

Citation for published version:

Buckley, BR, Thijssen, DHJ, Murphy, RC, Graves, L, Cochrane, M, Gillison, F, Crone, D, Wilson, P, Whyte, G & Watson, P 2020, 'Pragmatic evaluation of a co-produced physical activity referral scheme: A quasi-experimental study', *BMJ Open*, vol. 10, e034580. <https://doi.org/10.1136/bmjopen-2019-034580>

DOI:

[10.1136/bmjopen-2019-034580](https://doi.org/10.1136/bmjopen-2019-034580)

Publication date:

2020

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

BMJ Open

Pragmatic evaluation of a co-produced physical activity referral scheme: A quasi-experimental study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034580.R2
Article Type:	Original research
Date Submitted by the Author:	n/a
Complete List of Authors:	Buckley, Benjamin; Liverpool John Moores University, Physical Activity Exchange Thijssen, Dick; Liverpool John Moores University Murphy, Rebecca; Liverpool John Moores University, Research Institute for Sport and Exercise Sciences Graves, Lee; Liverpool John Moores University, Cochrane, Madeleine; Liverpool John Moores University, Physical Activity Exchange Gillison, Fiona; University of Bath, Department for Health Crone, Diane; Cardiff Metropolitan University Wilson, Philip; Brock University Whyte, Greg; Liverpool John Moores University Watson, Paula; Liverpool John Moores University, Research Institute for Sport and Exercise Sciences
Primary Subject Heading:	Sports and exercise medicine
Secondary Subject Heading:	Public health, Cardiovascular medicine
Keywords:	Cardiovascular Health, Self-Determination Theory, Exercise Referral, Behaviour Change, Translational Research

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1 Pragmatic evaluation of a co-produced physical activity referral scheme: A quasi-experimental
2 study

3 Benjamin J. R. Buckley^a, Dick H. J. Thijssen^{a,e}, Rebecca C. Murphy^a, Lee E. F. Graves^a,
4 Madeleine Cochrane^a, Fiona Gillison^b, Diane Crone^c, Philip M. Wilson^d, Greg Whyte^a and
5 Paula M. Watson^a

6
7 ^a Research Institute for Sport and Exercise Sciences, Liverpool John Moores University,
8 Liverpool, UK

9 ^b Department for Health, University of Bath, Bath, UK

10 ^c Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, UK

11 ^d Behavioural Health Sciences Research Lab, Department of Kinesiology, Brock University,
12 Ontario, Canada

13 ^e Radboud Institute for Health Sciences, Department of Physiology, Radboud University
14 Medical Center, Nijmegen, Netherlands

15
16
17 Correspondence to Dr Ben Buckley: B.J.Buckley@ljmu.ac.uk

18
19 Contributorship Statement

20 BJRB contributed to the study design, data collection, data analysis, and preparation of the final
21 document. PMW, DHJT, and RCM contributed to the study design, data analysis, and preparation of
22 the final document. MC contributed to the data collection and approved the final version. LEFG, FG,
23 DC, PW, and GW intellectually contributed to this paper and approved the final version.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Objectives. UK exercise referral schemes (ERSs) have been criticised for focusing too much on exercise prescription and not enough on sustainable physical activity (PA) behaviour change. Previously, a theoretically-grounded intervention (Co-PARS) was co-produced to support long-term PA behaviour change in individuals with health conditions. The purpose of this study was to investigate the effectiveness of Co-PARS compared to a usual care ERS and no treatment for increasing cardiorespiratory fitness.

Design. A three-arm quasi-experimental trial.

Setting. Two leisure centres providing a) Co-PARS, b) usual exercise referral care, and one no-treatment control.

Participants. 68 adults with lifestyle-related health conditions (e.g. cardiovascular, diabetes, depression) were recruited to Co-PARS, usual care, or no treatment.

Intervention. 16-weeks of physical activity behaviour change support delivered at 4, 8, 12, and 18 weeks, in addition to the usual care 12-week leisure centre access.

Outcome measures. Cardiorespiratory fitness, vascular health, PA, and mental wellbeing were measured at baseline, 12 weeks, and 6 months (PA and mental wellbeing only). Fitness centre engagement (Co-PARS and usual care) and behaviour change consultation attendance (Co-PARS) were assessed. Following an intention-to-treat approach, repeated-measures linear mixed models were used to explore intervention effects.

Results. Significant improvements in cardiorespiratory fitness ($p=.002$) and vascular health ($p=.002$) were found in Co-PARS compared to usual care and no-treatment at 12 weeks. No significant changes in PA or wellbeing at 12 weeks or 6 months were noted. Intervention engagement was higher in Co-PARS than usual care, though this was not statistically significant.

Conclusion. A co-produced PA behaviour change intervention led to promising improvements in cardiorespiratory and vascular health at 12 weeks, despite no effect for PA levels at 12 weeks or 6 months.

Trial registration: ClinicalTrials.gov: NCT03490747

Keywords: Cardiovascular Health; Self-Determination Theory; Exercise Referral; Behaviour Change Intervention; Translational Research.

Strengths and limitations of the study

- This study advances the literature on exercise referral effectiveness by pragmatically evaluating a co-produced physical activity referral intervention, which was underpinned by multiple stakeholders and behaviour change theory.
- The study documents the third phase of a novel and iterative approach which co-produced, piloted, and then evaluated (this study) a physical activity referral intervention that was deemed feasible to implement in practice.
- Objective and subjective measures provide insight into the potential effects for patient health.
- It is not possible to directly attribute intervention effects to the phased co-production approach, although supported by the Medical Research Council.
- A larger sample size is needed to substantiate findings.

Funding

This project was supported by a PhD studentship for Benjamin Buckley from Liverpool John Moores University. The 6-month data collection and analysis was supported by a financial grant from NHS Liverpool Clinical Commissioning Group.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BJRB contributed to the study design, data collection, data analysis, and preparation of the final document. PMW, DHJT, and RCM contributed to the study design, data analysis, and preparation of the final document. MC contributed to the data collection and approved the final version. LEFG, FG, DC, PW, and GW intellectually contributed to this paper and approved the final version.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Word count

~3000

Ethics approval and consent to participate

Full written consent was obtained from participants and the study was approved by NHS Research Ethics Committee (REC: 18/NW/0039 - Project: 238547).

Acknowledgements

1
2
3 1 We would like to thank the participants in this study for their time, the delivery staff and centre
4 2 managers for their ongoing support, and the initial development group involved in the co-production
5 3 process.
6 4
7 5

8
9
10 5 **INTRODUCTION**

11 6 Physical inactivity is the fourth leading cause of death worldwide and costs the UK an estimated £7.4
12 7 billion annually, including £0.9 billion to the NHS alone[1]. Exercise referral schemes (ERSs) provide a
13 8 promising framework to facilitate physical activity (PA) behaviour change in at-risk populations.
14 9 Typically, UK ERSs consist of a referral from a healthcare professional to a 12-16-week tailored exercise
15 10 programme provided by a qualified practitioner.

16 11 There is inconsistent evidence as to the effectiveness of ERSs on PA behaviour, mental well-being,
17 12 quality of life, and physical health outcomes [2–4]. More recently, however, promising effects of ERSs
18 13 have been demonstrated in Wales [5], Sweden [6], and Spain [7] and a systematic review identified
19 14 promising effects of UK ERSs on self-reported PA and cardiovascular health markers [8]. Prior and
20 15 colleagues [9] demonstrated that for every 11 participants referred to a 24-week ERS, 1 participant
21 16 went on to report achieving ≥90 min/week of PA at 12-months. For perspective, it is estimated that
22 17 67-167 patients (categorised as ≤10% cardiovascular disease (CVD) risk) need to receive statin
23 18 treatment for 5 years to prevent one major vascular event [10]. Whilst we are not suggesting PA
24 19 behaviour change is a comparable outcome to a serious clinical event, it is notable that replacing 30
25 20 minutes of TV viewing time with PA across the UK population, could reduce premature mortality by
26 21 5-15%, depending on activity intensity [11]. The majority of studies evaluating ERSs, however, have
27 22 drawn on self-reported PA data and future studies employing device-based measures are needed to
28 23 substantiate these observations.

29 24 Despite recent promise for the effectiveness of ERSs [7–9,12], substantial heterogeneity exists in both
30 25 design and delivery [13,14], reflecting varying assumptions on how best to promote health behaviour
31 26 change [15,16]. This limits potential scalability of ‘successful’ ERSs. Traditionally, ERSs have focussed
32 27 on short-term exercise prescription without appropriate evidence of effectiveness or underpinning of

behaviour change theory [17]. A recent attempt to integrate behaviour change theory into an ERS [18] however, showed no advantage over a standard ERS at 12 weeks or 6 months. The authors noted considerable implementation challenges when training staff, such as work-related demands that may have reduced the importance of the theory-based training. It is plausible that delivery staff asked to implement interventions designed by academics may lack ownership and feel less motivated/competent. One potential way to promote ownership and engagement might be to adopt a co-production approach, as a means of co-creating value across the public sector [19–21]. Though not a panacea, the involvement of practitioners, managers and service-users in co-production has potential to improve intervention relevance, fidelity, and effectiveness [22].

Previously, a theoretically-grounded PA referral scheme (Co-PARS) was co-produced by academics, policy-makers, practitioners, and service-users [23] in Liverpool, UK, with a focus on supporting sustainable PA behaviour change. Liverpool is the 3rd most deprived local authority in England and has the 2nd highest proportion of Lower Super Output Areas (LSOAs) in the most deprived 10% nationally [24]. Interventional work with at-risk patients is therefore critical and is aligned with the concept of proportionate universalism [25]. Underpinned by self-determination theory [24], the co-produced intervention differed from usual ERS care in its focus on PA behaviour change (rather than exercise prescription), and inclusion of frequent one-to-one consultations with exercise referral practitioners (compared to usual care which included formal contact at induction only). A pilot of Co-PARS [26] showed clinically meaningful improvements in cardiorespiratory fitness (CRF) and PA, although as we did not include a usual care control, it was unknown whether these effects were due to the fact participants were taking part in an ERS or due to the unique elements of Co-PARS. Furthermore, despite having very low CRF ($<27.7 \text{ ml.kg}^{-1}.\text{min}^{-1}$) [26] we found 64% of the baseline pilot sample were meeting the PA guidelines [27] of at least 150 minutes moderate-intensity PA per week (measured objectively via accelerometry). This suggested CRF may be a more appropriate primary outcome measure than PA for this low-fit population (whilst changing PA behaviour was the focus of the intervention, a target health outcome of this behaviour change was improved CRF). The pilot also

1
2
3 1 allowed the opportunity to investigate delivery processes, and we noted several areas that required
4
5 2 refinement in preparation for a controlled trial. These refinements included, increasing the number
6
7 3 of behaviour change consultations from four to five; enhanced focus on daily PA opportunities (rather
8
9 4 than focussing on activities offered at the fitness centre); adapting staff timetables to promote
10
11 5 consistency of care and to allow participant one-to-one consultations to take place in a private room;
12
13 6 and reducing practitioner paperwork. Building on our previous pilot work, the aim of the current study
14
15 7 was to investigate the effectiveness of Co-PARS compared to a usual care ERS and a no-treatment
16
17 8 control on change in cardiorespiratory fitness (CRF) at 12 weeks and PA and wellbeing at 6 months.
18
19

20
21 9 **METHODS**

22
23 10 **Study Design**

24
25
26 11 A three-arm quasi-experimental trial involving: 1. Co-PARS (delivered at fitness centre A); 2. usual care
27
28 12 ERS (delivered at fitness centre B); and 3. no-treatment control. This paper reports trial outcomes
29
30 13 (CRF, vascular health, PA, mental wellbeing) measured at baseline, 12 weeks, and 6 months (PA and
31
32 14 mental wellbeing only). Additional data were collected to investigate psychosocial processes of
33
34 15 change, intervention fidelity and cost-effectiveness; due to space limitations they are not considered
35
36 16 in the present manuscript, but findings can be obtained on request from p.m.watson@ljmu.ac.uk. Full
37
38 17 written consent was obtained from participants and the study was approved by NHS Research Ethics
39
40 18 Committee (REC: 18/NW/0039 - Project: 238547) and registered on ClinicalTrials.gov (NCT03490747).
41
42
43

44
45 19 **Patient and Public Involvement**

46
47 20 The intervention was previously co-produced, piloted, and adapted with substantial service user input
48
49 21 [23,26].
50
51

52 22 **Participants and Recruitment**

53
54 23 Inclusion criteria were the same for all three conditions (Co-PARS, usual care, no-treatment).
55
56 24 Participants were eligible if aged ≥ 18 years with a health-related risk factor (e.g. hypertension,
57
58 25 hyperglycaemia, obesity) and/or health condition (e.g. diabetes, cardiovascular disease, depression)
59
60

that may be alleviated by increasing PA levels. Participants with uncontrolled health conditions, severe psychological or neurological conditions were excluded. Participants for the Co-PARS and usual care arms were recruited from fitness centre A (Co-PARS) and fitness centre B (usual care) respectively (where they had been referred for exercise by a health professional). Reception staff at both centres provided study information and gained consent to pass participant details to the researcher. Participants for the no-treatment control were recruited via posters, electronic invitations, and email communications primarily at the university site. Participants were not eligible for the no-treatment control if they were currently attending an exercise referral scheme. Interested participants for all groups were sent an information sheet and baseline data collection was arranged.

Study Arms

Intervention arm components are presented in Figure 1.

Usual care exercise referral scheme (ERS – centre B). Usual care followed a standard ERS model of 12-week subsidised access to a fitness centre (swimming, gym, group classes). Participants met an exercise referral practitioner for an initial, 1-hour induction (week 1) during which a 12-week exercise programme was provided for the participant. Any further contact with a practitioner was informal and opportunistic. This system was already in place and was considered usual care for the local area. Centre B was chosen as a comparison centre due to its similarity in referral numbers and socio-economic make-up of the local population to centre A (where Co-PARS was being delivered). For example, based on areas within Liverpool ranked from 1 (most deprived) to 30 (least deprived), usual care ERS and Co-PARS were ranked respectively: 20th and 21st (income), 20th and 21st (employment), 22nd and 24th (Education) and 10th and 11th (living environment).

Co-produced PA referral scheme (Co-PARS – centre A)

Participants received the same 12-week subsidised access to a fitness centre as usual care plus a series of one-to-one behaviour change consultations (60-minute induction followed by 30-minute consultations at weeks 4, 8, 12 and 18). A log book was provided for each participant to set action

1
2
3 1 plans, log progress and facilitate consultation discussions. Consultations were delivered by exercise
4
5 2 referral practitioners in an autonomy supportive counselling style, drawing on the principles of self-
6
7 3 determination theory [28]. This additional support aimed to encourage habitual opportunities to
8
9 4 increase PA as well as activities available at the fitness centre. A full description of the theoretical
10
11 5 underpinning and behaviour change intervention components is available elsewhere [23].
12
13
14
15 6 Prior to the pilot of Co-PARS [26] practitioners received training in self-determination theory-based
16
17 7 communication strategies led by a sport and exercise psychologist (last author [PMW]), involving a
18
19 8 workshop, one-to-one sessions and follow-up group meetings. Following the pilot, a further series of
20
21 9 group meetings involving exercise referral practitioners and the research team were held to develop
22
23 10 aspects of delivery that required refinement (as outlined in the introduction). Full details of the
24
25 11 training are available from p.m.watson@ljmu.ac.uk.
26
27
28
29 12 **No-treatment control (NTC).** Participants received a lifestyle advice booklet only (offered to all study
30
31 13 arms at baseline data collection), based on national guidance for PA, nutrition, smoking cessation and
32
33 14 alcohol consumption.
34
35
36 15 [INSERT FIGURE 1 SOMEWHERE HERE]
37
38
39 16
40
41 17 **Outcome measures**
42
43
44 18 **Primary outcome:** *Cardio-respiratory fitness (CRF).* Maximal oxygen consumption ($\text{VO}_{2\text{max}}^2$) was
45
46 19 estimated via the sub-maximal Astrand-Rhyming cycle ergometer protocol [29]. The protocol is a
47
48 20 single-stage cycling test designed to elicit a steady-state heart rate over a period of ~6 minutes.
49
50
51 21 **Accelerometer-derived PA.** Tri-axial ActiGraph GT3x accelerometers (ActiGraph, Pensacola, FL, USA)
52
53 22 measured PA for 7 days, which have been validated in a comparable population [30]. Raw triaxial
54
55 23 acceleration values were converted into an omnidirectional measure of acceleration, referred to as
56
57 24 Euclidian norm minus one [31]. Minimum wear time was 10 hours per day and 3 days per week
58
59 25 including one weekend day [32]. The R package GGIR [31] facilitated extraction of user-defined

1 acceleration thresholds: 5.9 to 69.1 mg for light-intensity PA [33], 69.1 to 258.7 mg as moderate and
2 >258.7 mg as vigorous-intensity PA [34].

3 *Vascular health.* Our previous work has demonstrated carotid artery reactivity (CAR) may be a
4 promising outcome variable to assess in PA interventions for at-risk populations [35]. Further,
5 endothelial function may provide prognostic value beyond that of traditional risk factors [36] with an
6 increase of 1% in brachial artery flow-mediated dilation (FMD) associated with a 12-15% lower risk of
7 CV events [33,34]. FMD and CAR were measured using ultrasound techniques [35]. Both techniques
8 measure vascular endothelial function and have independently predicted future risk of cardiovascular
9 events in humans [36,37]. Blood pressure was measured in the supine position using an automated
10 blood pressure device (Omron Healthcare UK Limited, Milton Keynes, UK).

11 *Anthropometric measures.* Since obesity is a critical risk factor for poor health and cardiovascular
12 disease, anthropometric variables were measured to investigate potential intervention effects on
13 body mass. Waist-to-height ratio is a stronger predictor of early health risk than Body Mass Index
14 (BMI) alone [38], therefore we collected both BMI (mass in kg / stature in m²) and waist-to-height
15 ratio (waist circumference / stature).

16 *Mental wellbeing.* As PA is known to enhance mental wellbeing [39] and clinical populations are more
17 susceptible to mental ill-health [40], it was important to identify whether Co-PARS led to any changes
18 in mental health (positive or negative). Mental wellbeing was measured using the 14-item Warwick-
19 Edinburgh Mental Well-being Scale (WEMWBS; [41], which asks participants to rate their
20 psychological wellbeing (e.g. "I've been feeling cheerful") over the previous 2 weeks (measured on a
21 likert scale of 1 (none of the time) to 5 (all of the time)).

22 *Fitness centre engagement (Co-PARS and usual care only).* The number of occasions participants
23 attended the fitness centre between baseline and 12 weeks (weekly attendance) and 12 weeks to 6
24 months (monthly attendance) was obtained from computerised attendance records. When
25 measuring intervention engagement it was deemed inappropriate to calculate the mean number of

1 sessions per week, since this could exaggerate the engagement of individuals who attended with
2 high frequency in the early weeks then dropped out (when compared with individuals who attended
3 moderately but consistently for the full 12 weeks). Therefore a formula was used to calculate a
4 percentage for ‘12-week engagement’ (based on the recommended bi-weekly attendance):

$$\left(\frac{((n1*0.5) + (n2) + (n3*1.2))}{12} \right) * 100$$

n1 = number of weeks in which participant attends once only
n2 = number of weeks in which participant attends twice
n3 = number of weeks in which participant attends three or more times

10 This formula took into account both *frequency* and *consistency* of attendance to yield a percentage
11 score that ranged from 0% (no attendance) to 120% (attendance of three or more times per week
12 for the whole 12 weeks).

13 Monthly attendance post-12 weeks was calculated as a mean attendance across months 4 to 6,
14 therefore did not take consistency of attendance into account.

15 *Behaviour change consultation attendance (Co-PARS only).* The number of consultations offered and
16 attended were measured by exercise referral practitioners at induction, 4, 8, 12, and 18 weeks.

17 **Sample size**

18 Sample size was determined to detect a meaningful difference in CRF at 12 weeks based on our pilot
19 results [26]. To detect a difference of 2 ml.kg⁻¹min⁻¹ between Co-PARS and usual care, 42 participants
20 were required per arm (f= .25, p= .05, power = .80). To detect a difference of 3.2 ml.kg⁻¹min⁻¹ between
21 the intervention arms and the no-treatment control, 17 participants were required for the no-
22 treatment control (f= .5, p= .05, power = .80). Thus, a total sample of 101 participants were required.

23 **Statistical analyses**

24 An intention-to-treat approach was used assuming no change in non-respondents (last observation
25 carried forward) to produce a conservative estimate of intervention effects. Delta changes (Δ) from

pre- to post-intervention were calculated for each group and entered as the dependent variable in repeated measures linear mixed model analyses. A random intercept model was used with fixed effects for study arm (Co-PARS, usual care ERS, no-treatment control) and time (baseline-to-week-12 change, week-12-to-6-month change, and baseline-to-6-month change) and participants included as random effects. Least squared difference (LSD) was used for post hoc testing. Testing for baseline differences to identify covariates was avoided, as this method has been demonstrated to inflate bias, instead pre-intervention was entered into the model as a covariate. Furthermore, all linear mixed model analyses were repeated with age and employment as covariates as a comparison to the results presented in this study (with baseline score as a covariate) due to their known prognostic value. Using age and employment as covariates resulted in no change in inferences presented in this study. One-way ANOVAs were used to compare baseline values between intervention arms. Fitness centre engagement was determined as described above. Behaviour change consultation attendance is presented descriptively. For non-normally distributed data, median and interquartile range is presented and within group median change was calculated via Wilcoxon signed-rank tests.

RESULTS

Participants. 68 participants provided baseline data, 56 of whom provided 12-week data, and 58 of whom provided 6-month data (figure 2).

Baseline characteristics (table 1). No significant differences were noted between arms for age, sex, ethnicity, BMI, referral reason, or accelerometer-derived PA levels ($p>.05$). Full-time employment status ($p=.001$) and CRF ($p=.015$) were significantly higher in the control compared to usual care and Co-PARS. Smoking status was significantly higher in usual care compared to Co-PARS and control ($p=.010$). Mental wellbeing was significantly lower in Co-PARS compared to control ($p=.023$).

[INSERT FIGURE 2 SOMEWHERE HERE]

Table 1. Baseline characteristics presented as Mean ± SD or % (n) of group.

	Co-produced PA referral	Usual care ERS	No-treatment control	Between arm
	(n=33)	(n=19)	(n=16)	p-value
Age (years)	57 ± 12	53 ± 16	48 ± 15	p=.319
Female (% of sample)	58 (19)	47 (9)	56 (9)	p=.774
White British (% of sample)	82 (27)	95 (18)	75 (12)	p=.132
Full-time employment (% of sample)	18 (6)	26 (5)	62 (10)	p=.001
Never smoked (% of sample)	73 (24)	37 (7)	81 (13)	p=.002
Body mass index (kg/m²)	31 ± 7	33 ± 6	29 ± 6	p=.226
Systolic blood pressure (mmHg)	131 ± 11	138 ± 18	123 ± 12	p=.010
Primary referral reason / health concern (control)				p=.132
Cardiometabolic (% of sample)	67 (22)	43 (8)	62 (10)	-
Cancer (% of sample)	6 (2)	5 (1)	6 (1)	-
Mental Health (% of sample)	18 (6)	26 (5)	19 (3)	-
Musculoskeletal (% of sample)	9 (3)	26 (5)	13 (2)	-
Comorbidity (% of sample)	85 (28)	100 (19)	81 (13)	p=.166
Meeting the PA guidelines (% of sample)*	73 (22)	71 (10)	93 (13)	p=.223

P-values represent between arm baseline effects. There was no between arm effect for referral reason, thus no between arm *p-values* are provided for referral reason sub groups.

*Chief Medical Officers’ 2019 physical activity guidelines: 150 minutes of moderate-intensity physical activity per week.

Baseline-to-12-Week effects

Raw outcome values are presented for baseline, week 12, and 6 months in Table 2. There was a significant effect for study arm in baseline-to-12-week change in CRF (*p*=.002). Post hoc testing

revealed a significantly higher CRF change in Co-PARS (2.4) compared to the ERS (0.3; $p=.021$) and control (-0.6; $p=.001$), but no difference between the ERS and control ($p=.314$). A significant effect for study arm was found in change in FMD% ($p=.002$), with FMD% change significantly higher in Co-PARS (2.4) compared to control (-1.1; $p=.001$) but not the ERS (0.8; $p=.099$). The change in FMD% was not significantly different between the ERS and control ($p=.71$). No statistically significant study arm effects were noted for changes in CAR%, blood pressure, resting heart rate, anthropometric measures, PA or WEMWBS at 12 weeks ($p>.05$).

Baseline-to-6-month effects

No statistically significant study arm effects were noted for change in WEMWBS or PA at 6 months ($p>.05$).

Fitness centre engagement (Co-PARS and usual care ERS) and consultation attendance (Co-PARS only).

Table 3 reports the participant fitness centre engagement data for the Co-PARS and usual care ERS. Although not statistically significant, Co-PARS engagement was 9% higher, participants attended the fitness centre on average 3 times more per month, and 23% more participants were attending the fitness centre beyond 6-months follow-up compared to usual care. Co-PARS behaviour change consultation attendance is reported in Table 4.

Table 2. Cardiometabolic health outcomes and PA levels at baseline, 12 weeks, 6 months, and between arm baseline-to 12-week or 6-month effect. All variables are presented as Mean ± SD.

	Co-PARS			Usual Care ERS			No-Treatment Control			Between arm effect <i>p</i> -value ^(a)
	Baseline	Week 12	6 Month	Baseline	Week 12	6 Month	Baseline	Week 12	6 Month	
<i>Fitness</i> (n=56)										
<i>CRF</i> <small>ml.kg.⁻¹min⁻¹</small>	22.2±7	24.6±7	-	23.3±6.6	23.6±7	-	29.6±9.2	28.9±8.7	-	<i>p</i> =.002
<i>Physical Activity</i>										
<i>GT3x</i> (n= 61) <small>Mins.day</small>										
<i>Light intensity</i>	90±52	98±64	107±75	98±36	93±31	158±145	90±37	101±33	86±40	<i>p</i> =.332
<i>Moderate intensity</i>	44±32	42±29	42±33	43±28	43±30	55±55	60±31	65±24	54±21	<i>p</i> =.260
<i>Vigorous intensity</i>	1±3	1±2	1±2	1±2	1±1	1±2	2±4	2±3	3±8	<i>p</i> =.108
<i>Vascular Ultrasound</i> (n=64)										
<i>CAR%</i>	1.7±2.7	2.8±2.2	-	2.7±1.8	3.9±2.8	-	2.5±2.7	1.7±2.7	-	<i>p</i> =.073
<i>CAR Baseline</i> <small>cm</small>	0.69±0.07	0.69±0.06	-	0.69±0.08	0.7±0.09	-	0.65±0.07	0.64±0.06	-	<i>p</i> =.130
<i>FMD%</i>	4.4±2.3	6.8±2.7	-	4.2±2	5±2.1	-	6.2±2.1	5.2±2.8	-	<i>p</i> =.002
<i>FMD Baseline</i> <small>cm</small>	0.39±0.07	0.38±0.06	-	0.39±0.09	0.41 0.08	-	0.38±0.08	0.37±0.06	-	<i>p</i> =.728
<i>Cardiometabolic</i> (n=68)										
<i>BMI</i> <small>kg.m2</small>	31±7	30±7	-	33±6	32±6	-	29±6	29±6	-	<i>p</i> =.323
<i>WHR</i>	62±9	61±10	-	64±8	63±8	-	56±9	56±9	-	<i>p</i> =.261
<i>SBP</i> <small>mmHg</small>	131±11	127±12	-	138±18	132±15	-	123±12	118±13	-	<i>p</i> =.937
<i>DBP</i> <small>mmHg</small>	73±7	71±8	-	73±9	71±11	-	72±11	68±10	-	<i>p</i> =.584
<i>RHR</i> <small>bpm</small>	70±10	65±10		70±12	68±11		66±12	63±9		<i>p</i> =.540
<i>Mental Wellbeing</i> (n=68)										
WEMWBS	46±9	51±10	48±10	49±10	52±11	50±13	53±9	56±9	53±10	<i>p</i> =.796

Co-PARS, Co-produced PA referral scheme; ERS, Exercise referral scheme; CRF, Cardiorespiratory Fitness; GT3x, Accelerometer; CAR, Carotid artery reactivity; FMD, Flow-mediated dilation; BMI, Body Mass Index; WHR, Waist-to-Height ratio; SBP, Systolic blood pressure; DBP, Diastolic blood pressure; RHR, Resting heart rate, WEMWBS, Warwick-Edinburgh Mental Wellbeing Scale

^a *F*-statistic for between arm baseline-to-6-month change or baseline-to-week 12 change if variable not collected at 6 months.

Missing data was due to inability to complete the CRF test (n=12), inability to complete the vascular ultrasound protocols (n=4), and insufficient accelerometer wear time or non-return (n=7).

Table 3. Fitness centre engagement.

	Co-PARS (n=33)	Usual Care (n=19)	Between centre difference
% Engagement ^a (Mean ± SD)	42±29	33±27	<i>p</i> =.267
Number of fitness centre visits (per person per month) week 12 to 6 months (Med, IQR)	3(0-14)	0 (0-1)	<i>p</i> =.072
% of baseline sample who attended fitness centre at least once beyond 6 months (% of sample, n)	39 (13)	16 (3)	<i>p</i> =.101

^aBased on the formula $((n1*0.5)+(n2)+(n3*1.2))/12 * 100$; n1=number of weeks in which participant attends once only; n2=number of weeks in which participant attends twice; n3=number of weeks in which participant attends three or more times. ^a**Engagement**; based on a recommended attendance of twice weekly, a formula was used to calculate a percentage for "12-week engagement", which took into account both frequency and consistency of attendance (see methods).

Table 4. Co-PARS behaviour change consultation attendance (based on baseline sample of 33 participants).

Consultation	% Booked (n)	% Attended (n)
Induction	91(30)	93(28)
Week 4	82(27)	78(21)
Week 8	67(22)	91(20)
Week 12	64(21)	81(17)
Week 18	55(18)	50(9)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

This was the first study to investigate the effectiveness of a theoretically-grounded, co-produced PA referral scheme (Co-PARS) compared to a usual care ERS and no treatment. Despite challenges in recruitment that meant the study was statistically underpowered, the findings demonstrated significant and clinically meaningful improvements in CRF and vascular health in Co-PARS compared to the usual care and no treatment. No statistically significant effects were noted for accelerometer-derived PA levels or mental wellbeing at 12-weeks or 6-months.

The effect of usual care ERSs compared to theoretically-grounded interventions on CRF has not been previously explored. We observed a significant increase in CRF in Co-PARS compared to usual care and a no-treatment control. According to values reported by Clausen *et al.* [42] both Co-PARS (22 ml.kg.⁻¹min⁻¹) and usual care (23 ml.kg.⁻¹min⁻¹) participants were below the lower limit of ‘healthy’ (27.7 ml.kg.⁻¹min⁻¹) for baseline CRF [43]. As low CRF is associated with a substantially elevated risk of all-cause mortality [43], the magnitude of change demonstrated in Co-PARS (2.4 ml.kg.⁻¹min⁻¹) may be clinically meaningful. For example, in at-risk populations, relatively small magnitudes (≤1 ml.kg.⁻¹min⁻¹) have been shown to significantly reduce clustered cardiometabolic risk [44]. Thus, Co-PARS was effective at improving CRF in individuals with low CRF by a clinically meaningful amount.

Promising improvements in vascular health were also noted in the Co-PARS group, with brachial artery FMD significantly improved compared to usual care and control arms. Although CAR was not statistically different between arms, both Co-PARS and usual care demonstrated a potentially meaningful within-arm improvement compared with no treatment, which exhibited a deterioration in vascular health. Such improvements in vascular measures may have prognostic implications. For example, a 1% increase in FMD has been suggested to reduce the future risk of CVD events by 13% [36].

Despite low baseline CRF, a substantial percentage of Co-PARS (73%) and usual care (71%) participants were meeting the Department of Health [45] guidelines of 150 minutes of moderate-intensity PA per

1 week. We observed a similar finding in our pilot [26] and subsequently raised the question as to the use of PA guidelines to assess eligibility for ERSs (NICE, 2014), as it appears from our data that individuals classified as “physically active” can still be very unfit and therefore can benefit from ERSs in terms of improved fitness and cardiometabolic health. A further discrepancy was noted in the lack of change in PA levels in Co-PARS, despite improved CRF. It is possible measurement issues contributed to this discrepancy. Accelerometers can measure certain types of PA such as walking, running, and stair climbing [46]. They may not, however, sufficiently identify activities typical of an ERS delivered within a fitness centre environment (e.g. cycling, resistance training, circuits, swimming). Given Co-PARS had higher (albeit non-significant) fitness centre engagement compared to usual care, it is possible PA changes occurred that were not detected by the accelerometry data. Consideration therefore needs to be given to the appropriateness of accelerometers to measure PA in ERSs. Alternative methods such as heart-rate monitors combined with self-report data may be worthy of consideration, although further work would be required to develop standardized data collection and analysis protocols (taking into account the limitations of each of these methods if used in isolation [47]). Researchers are therefore urged to consider CRF as a primary outcome in ERSs until appropriate alternative methods of measuring PA behaviour are developed. Ultimately, it is not clear why the increase in fitness occurred without a corresponding change in PA and further research is required to elucidate the relationship between PA and fitness in this population.

In addition to physiological health outcomes, we found baseline mental wellbeing to be below the national average (score of 50) in both Co-PARS (46) and usual care (49), but not the control (53) [48]. Despite no significant between-group effect for mental wellbeing, within-group changes at 12 weeks were deemed clinically meaningful for Co-PARS (5) and usual care (3) but not in the no treatment control. It is notable that the post-intervention magnitude of change observed in mental wellbeing for Co-PARS (5) was larger than that observed in a meta-analysis encompassing >23,000 participants across 13 different ERSs (3), which were comparable in nature to the usual care ERS in this study [49].

1
2
3 1 From the 6-month data it appeared the scheme was not effective at promoting *sustained* PA behaviour
4
5 2 change or mental wellbeing improvements. It must be noted, however, that the wellbeing levels were
6
7 3 still higher than baseline and even small magnitudes of change (1-3) may be meaningful in clinical
8
9 4 populations [50]. As discussed earlier, it may be that measuring PA using the methods described in
10
11 5 this study prevented the identification of activities typical of a fitness centre environment. This notion
12
13 6 is supported by the post-week-12 attendance data, which highlighted Co-PARS participants were
14
15 7 regularly attending the fitness centre whereas the usual care participants were not. Challenges of
16
17 8 maintaining sustained health outcomes post-ERSs have been highlighted elsewhere [3]. And whilst a
18
19 9 recent systematic review reported longer length schemes (>20 weeks) may be more effective than
20
21 10 shorter schemes [8], the four long ERSs (20-26 weeks) collected pre-post data only. Thus we do not
22
23 11 know if longer length ERSs result in enhanced health outcomes *post intervention* compared with
24
25 12 shorter schemes. To determine if longer length schemes are indeed more effective, longer-term
26
27 13 follow-up data collection is required, ideally at 6 and 12 months post intervention [51].
28
29
30
31
32
33 14 Through a phased approach we have assessed the effectiveness of Co-PARS resulting from several
34
35 15 years of co-production. Whilst the effects of co-production are difficult to isolate, a comparison of
36
37 16 results at different stages of intervention refinement suggests the phased development approach had
38
39 17 some positive effects. Unpublished engagement data from centre A in 2014-2015 (when the centre
40
41 18 was running a usual care ERS) shows that engagement improved after the introduction of Co-PARS
42
43 19 (42% vs 28% in 2014-2015), whereas engagement reduced in the usual care centre over the same
44
45 20 period (32% vs 37% in 2014-2015). Furthermore, consultation attendance for Co-PARS in the current
46
47 21 study was substantially higher than in our previous pilot (54% attended induction plus ≥ 3 behaviour
48
49 22 change consultations, vs 9% in the pilot [26]), which may have been a reflection of refinements made
50
51 23 to the intervention after the pilot (e.g. improved focus on holistic PA, improved monitoring
52
53 24 procedures, improved continuity of instructors). These improvements in engagement highlight the
54
55 25 importance of allowing time for complex interventions to develop [52], and are particularly promising
56
57 26 given the effectiveness of ERSs are highly dependent on participant adherence [5,21]. Furthermore,
58
59
60

1 this study has demonstrated how investing in the “bottom-up” development of an intervention can
2 lead to an effective and sustainable model. We therefore support the arguments of Rutter and
3 colleagues [53] in that a shift in thinking is needed, instead of asking whether an intervention works
4 to fix a problem, researchers should aim to identify if and how it contributes to reshaping a system in
5 a favourable way. As such, we propose the co-production and implementation process may be as
6 important as the scheme content itself.

7 **Methodological considerations**

8 This is the first known study to investigate the effectiveness of a co-produced PA referral scheme (Co-
9 PARS) in comparison to usual care and a no-treatment control. Our novel approach addresses an
10 important gap in the sport and exercise medicine literature [54], in that we employed rigorous
11 laboratory-based instruments to measure health outcomes that can be achieved through an
12 ecologically valid, “real-world” intervention. We observed a very high retention at 6-month follow up,
13 which may be due in part to the fact many of the participants were retired (and therefore may have
14 more available time). It is possible also that the high retention was facilitated by the co-production
15 process, which involved ongoing relationships between the research and delivery teams (and
16 therefore helped with the logistics of returning accelerometers for the co-PARS and usual care
17 groups). Whilst this paper highlights many strengths of co-production, we do not wish to present co-
18 production as a panacea [19] and it is important potential challenges and costs are considered prior
19 to undertaking such an approach [21,22].

20 We must acknowledge some limitations of the study. Whilst there is a need for high-quality RCTs of
21 theoretically informed approaches to PA behaviour change [3], several pragmatic reasons meant an
22 RCT approach was not appropriate for the present study. Firstly, it was important participants could
23 choose the most convenient fitness centre. Secondly, it was important we continued work with the
24 same fitness centre and staff (following co-production [23] and pilot [26] phases) in order to develop
25 the intervention to the point where it was deemed to have a worthwhile effect [52]. A pragmatic

1
2
3 1 research approach was therefore deemed most appropriate to evaluate Co-PARS with high ecological
4
5 2 validity. Pragmatic constraints (e.g. fitness centre refurbishments, staff illness) did however mean the
6
7 3 required sample size was not achieved, thus inferences of effectiveness need to be taken with caution.
8
9
10 4 This is particularly true for the PA data, where the relatively high variability (compared with CRF) may
11
12 5 have contributed to the lack of change observed in PA in this study. It is recommended future work
13
14 6 considers pragmatic risks and contingencies when planning recruitment and plans sufficient time to
15
16 7 cope with recruitment delays. For pragmatic reasons, not all outcomes were collected at 6-months
17
18 8 follow-up and further research is needed to collect long-term, objective health data following PA
19
20 9 referral schemes. Finally, it must be noted that while the trial registration appears to be retrospective
21
22
23 10 (April 6th 2018), the initial submission was several months prior to this (January 11th 2018). Final sign-
24
25 11 off was delayed due to capacity issues within the research team.
26
27
28

29 12 **CONCLUSION**

30
31
32 13 A co-produced, theoretically-grounded PA referral scheme (Co-PARS) led to improved CRF and
33
34 14 vascular health in at-risk individuals when compared to usual care and no treatment. In addition,
35
36 15 clinically meaningful improvements in vascular health and mental wellbeing were observed at 12-
37
38 16 weeks in both Co-PARS and usual care, but not the no treatment control group. Of note, PA remained
39
40 17 unchanged at 12-weeks and 6-months follow-up. Adopting a phased approach has enabled multi-
41
42 18 stakeholder input and ongoing intervention refinement, resulting in an intervention that showed
43
44 19 promising effects on engagement and clinically meaningful improvements to participant health.
45
46
47
48
49 20

50
51
52
53
54
55
56
57
58
59
60

Figure Legends

Figure 1. 'PaT Plot' describing intervention arm components.[55]

Figure 2. Participant flow diagram within the three study arms (March 2018-January 2019).

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1 Public Health England. Physical activity: applying All Our Health. 2019.

2 Dugdill L, Graham RC, McNair F. Exercise referral: the public health panacea for physical activity promotion? A critical perspective of exercise referral schemes; their development and evaluation. *Ergonomics* 2005;**48**:1390–410. doi:10.1080/00140130500101544

3 Pavey T, Taylor A, Fox K, *et al.* Effect of exercise referral schemes in primary care on physical activity and improving health outcomes: systematic review and meta-analysis. *Bmj* 2011;**343**:d6462. doi:10.1136/bmj.d6462

4 Pavey T, Anokye N, Taylor A, *et al.* The clinical effectiveness and cost-effectiveness of exercise referral schemes: a systematic review and economic evaluation. *Health technology assessment (Winchester, England)* 2011;**15**:i–xii, 1–254. doi:10.3310/hta15440

5 Murphy SM, Edwards RT, Williams N, *et al.* An evaluation of the effectiveness and cost effectiveness of the National Exercise Referral Scheme in Wales, UK: a randomised controlled trial of a public health policy initiative. *Journal of epidemiology and community health* 2012;**66**:745–53. doi:10.1136/jech-2011-200689

6 Onerup A, Arvidsson D, Blomqvist Å, *et al.* Physical activity on prescription in accordance with the Swedish model increases physical activity: a systematic review. *Br J Sports Med* 2018;;bjsports-2018-099598. doi:10.1136/bjsports-2018-099598

7 Martín-Borràs C, Giné-Garriga M, Puig-Ribera A, *et al.* A new model of exercise referral scheme in primary care: is the effect on adherence to physical activity sustainable in the long term? A 15-

- 1 month randomised controlled trial. *BMJ Open* 2018;**8**:e017211. doi:10.1136/bmjopen-2017-017211
- 2
- 3
- 4
- 5
- 6
- 7
- 8 8 Rowley N, Mann S, Steele J, *et al.* The effects of exercise referral schemes in the United Kingdom in
- 9 those with cardiovascular, mental health, and musculoskeletal disorders: a preliminary systematic
- 10 review. *BMC Public Health* 2018;**18**:949. doi:10.1186/s12889-018-5868-9
- 11
- 12
- 13
- 14
- 15
- 16
- 17 9 Prior F, Coffey M, Robins A, *et al.* Long-Term Health Outcomes Associated With an Exercise
- 18 Referral Scheme: An Observational Longitudinal Follow-Up Study. *Journal of Physical Activity and*
- 19 *Health* 2019;**1**:1–6. doi:10.1123/jpah.2018-0442
- 20
- 21
- 22
- 23
- 24
- 25
- 26 10 Taylor F, Huffman MD, Macedo A, *et al.* Statins for the primary prevention of cardiovascular
- 27 disease. *Cochrane Db Syst Rev* 2013;**1**:CD004816. doi:10.1002/14651858.cd004816.pub5
- 28
- 29
- 30
- 31
- 32
- 33 11 Wijndaele K, Sharp SJ, Wareham NJ, *et al.* Mortality Risk Reductions from Substituting Screen
- 34 Time by Discretionary Activities. *Med Sci Sport Exer* 2017;**49**:1111–9.
- 35
- 36
- 37 16 doi:10.1249/mss.0000000000001206
- 38
- 39
- 40
- 41
- 42 12 Craike M, Wiesner G, Enticott J, *et al.* Equity of a government subsidised exercise referral scheme:
- 43 A population study. *Social Science & Medicine* Published Online First: 2018.
- 44
- 45
- 46 20 doi:10.1016/j.socscimed.2018.09.023
- 47
- 48
- 49
- 50
- 51 13 Craig A, Dinan S, Smith A, *et al.* Exercise Referral Systems: A National Quality Assurance
- 52 Framework. *Department of Health: London* Published Online First: 2001.discovery.ucl.ac.uk
- 53
- 54
- 55
- 56
- 57 14 Pavey T, Taylor A, Hillsdon M, *et al.* Levels and predictors of exercise referral scheme uptake and
- 58 adherence: a systematic review. *J Epidemiol Commun H* 2012;**66**:737–44. doi:10.1136/jech-2011-
- 59
- 60

1
2
3 1 200354
4
5 2
6
7
8 3 15 Littlecott HJ, Moore GF, Moore L, *et al.* Psychosocial mediators of change in physical activity in
9
10 4 the Welsh national exercise referral scheme: secondary analysis of a randomised controlled trial.
11
12 5 *International Journal of Behavioral Nutrition and Physical Activity* 2014;**11**:1–11.
13
14 6 doi:10.1186/s12966-014-0109-9
15
16 7
17
18 8 16 Hanson CL, Oliver EJ, Dodd-Reynolds CJ, *et al.* How do participant experiences and characteristics
19
20 9 influence engagement in exercise referral? A qualitative longitudinal study of a scheme in
21
22 10 Northumberland, UK. *Bmj Open* 2019;**9**:e024370. doi:10.1136/bmjopen-2018-024370
23
24 11
25
26 12 17 Sowden S, Raine R. Running along parallel lines: how political reality impedes the evaluation of
27
28 13 public health interventions. A case study of exercise referral schemes in England. *Journal of*
29
30 14 *epidemiology and community health* 2008;**62**:835–841. doi:10.1136/jech.2007.069781
31
32 15
33
34 16 18 Duda JL, Williams GC, Ntoumanis N, *et al.* Effects of a standard provision versus an autonomy
35
36 17 supportive exercise referral programme on physical activity, quality of life and well-being indicators:
37
38 18 a cluster randomised controlled trial. *The international journal of behavioral nutrition and physical*
39
40 19 *activity* 2014;**11**:10. doi:10.1186/1479-5868-11-10
41
42 20
43
44 21 19 Ostrom E. Crossing the great divide: Coproduction, synergy, and development. *World*
45
46 22 *Development* 1996;**24**:1073–87. doi:10.1016/0305-750x(96)00023-x
47
48 23
49
50 24 20 Clarke D, Jones F, Harris R, *et al.* What outcomes are associated with developing and
51
52 25 implementing co-produced interventions in acute healthcare settings? A rapid evidence synthesis.
53
54 26 *BMJ open* 2017;**7**:e014650. doi:10.1136/bmjopen-2016-014650
55
56
57
58
59
60

- 1
- 2
- 3 1
- 4
- 5 2 21 Farrance C, Tsofliou F, Clark C. Adherence to community based group exercise interventions for
- 6
- 7
- 8 3 older people: A mixed-methods systematic review. *Preventive medicine* 2016;**87**:155–66.
- 9
- 10 4 doi:10.1016/j.ypmed.2016.02.037
- 11
- 12 5
- 13
- 14 6 22 Rycroft-Malone J, Burton CR, Bucknall T, *et al.* Collaboration and Co-Production of Knowledge in
- 15
- 16 7 Healthcare: Opportunities and Challenges. *International journal of health policy and management*
- 17
- 18 8 2016;**5**:221–3. doi:10.15171/ijhpm.2016.08
- 19
- 20
- 21 9
- 22
- 23 10 23 Buckley B, Thijssen D, Murphy R, *et al.* Making a move in exercise referral: co-development of a
- 24
- 25 11 physical activity referral scheme. *Journal of Public Health* Published Online First: 2018.
- 26
- 27 12 doi:10.1093/pubmed/fdy072
- 28
- 29
- 30 13
- 31
- 32 14 24 The English indices of deprivation 2019. 2019.
- 33
- 34 15
- 35
- 36 16 25 Carey G, Crammond B, Leeuw E. Towards health equity: a framework for the application of
- 37
- 38 17 proportionate universalism. *Int J Equity Health* 2015;**14**:81. doi:10.1186/s12939-015-0207-6
- 39
- 40 18
- 41
- 42 19 26 Buckley BJ, Thijssen DH, Murphy RC, *et al.* Preliminary effects and acceptability of a co-produced
- 43
- 44 20 physical activity referral intervention. *Health Educ J* 2019;:001789691985332.
- 45
- 46 21 doi:10.1177/0017896919853322
- 47
- 48 22
- 49
- 50
- 51 23 27 Department of Health & Social Care. UK Chief Medical Officers' Physical Activity Guidelines. 2019.
- 52
- 53 24
- 54
- 55
- 56 25 28 Ryan R, Deci E. Self-determination theory and the facilitation of intrinsic motivation, social
- 57
- 58 26 development, and well-being. *The American psychologist* 2000;**55**:68–78.
- 59
- 60

1
2
3 1
4
5 2 29 Astrand I. Aerobic work capacity in men and women with special reference to age. *Acta*
6
7 3 *physiologica Scandinavica Supplementum* 1960;**49**:1–92.
8
9 4
10
11 5 30 Kelly LA, McMillan DG, Anderson A, *et al.* Validity of actigraphs uniaxial and triaxial
12
13 6 accelerometers for assessment of physical activity in adults in laboratory conditions. *BMC Medical*
14
15 7 *Physics* 2013;**13**:1–7. doi:10.1186/1756-6649-13-5
16
17 8
18
19 9 31 Hees VT, Gorzelniak L, Leon E, *et al.* Separating movement and gravity components in an
20
21 10 acceleration signal and implications for the assessment of human daily physical activity. *PLoS one*
22
23 11 2013;**8**:e61691. doi:10.1371/journal.pone.0061691
24
25 12
26
27 13 32 Matthews CE, Hagströmer M, Pober DM, *et al.* Best Practices for Using Physical Activity Monitors
28
29 14 in Population-Based Research. *Medicine & Science in Sports & Exercise* 2012;**44**:S68.
30
31 15 doi:10.1249/MSS.0b013e3182399e5b
32
33 16
34
35 17 33 Bakrania K, Yates T, Rowlands AV, *et al.* Intensity Thresholds on Raw Acceleration Data: Euclidean
36
37 18 Norm Minus One (ENMO) and Mean Amplitude Deviation (MAD) Approaches. *PLOS ONE*
38
39 19 2016;**11**:e0164045. doi:10.1371/journal.pone.0164045
40
41 20
42
43 21 34 Hilded M, Hees Vt, Hansen B, *et al.* Age Group Comparability of Raw Accelerometer Output from
44
45 22 Wrist- and Hip-Worn Monitors. *Medicine & Science in Sports & Exercise* 2014;**46**:1816.
46
47 23 doi:10.1249/mss.0000000000000289
48
49 24
50
51 25 35 Buckley B, Watson PM, Murphy RC, *et al.* Carotid artery function is restored in subjects with
52
53 26 elevated cardiovascular disease risk following a 12-week physical activity intervention. *Canadian*
54
55
56
57
58
59
60

- 1 *Journal of Cardiology* Published Online First: 2018. doi:10.1016/j.cjca.2018.10.015
- 2
- 3
- 4
- 5
- 6
- 7
- 8 36 Inaba Y, Chen JA, Bergmann SR. Prediction of future cardiovascular outcomes by flow-mediated
- 9
- 10 vasodilatation of brachial artery: a meta-analysis. *The international journal of cardiovascular imaging*
- 11
- 12 2010;**26**:631–40. doi:10.1007/s10554-010-9616-1
- 13
- 14
- 15
- 16
- 17 37 van Mil A, Pouwels S, Wilbrink J, *et al.* Carotid Artery Reactivity Predicts Events in Peripheral
- 18
- 19 Arterial Disease Patients. *Annals of Surgery* 2017. doi:10.1097/SLA.0000000000002558
- 20
- 21
- 22
- 23
- 24 38 Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist
- 25
- 26 circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis.
- 27
- 28 *Obesity reviews : an official journal of the International Association for the Study of Obesity*
- 29
- 30 2012;**13**:275–86. doi:10.1111/j.1467-789X.2011.00952.x
- 31
- 32
- 33
- 34
- 35 39 Paluska SA, Schwenk TL. Physical Activity and Mental Health. *Sports Med* 2000;**29**:167–80.
- 36
- 37 doi:10.2165/00007256-200029030-00003
- 38
- 39
- 40
- 41
- 42 40 Barnett K, Mercer SW, Norbury M, *et al.* Epidemiology of multimorbidity and implications for
- 43
- 44 health care, research, and medical education: a cross-sectional study. *Lancet* 2012;**380**:37–43.
- 45
- 46 doi:10.1016/s0140-6736(12)60240-2
- 47
- 48
- 49
- 50
- 51 41 Tennant R, Hiller L, Fishwick R, *et al.* The Warwick-Edinburgh Mental Well-being Scale
- 52
- 53 (WEMWBS): development and UK validation. *Health Qual Life Out* 2007;**5**:1–13. doi:10.1186/1477-
- 54
- 55 7525-5-63
- 56
- 57
- 58
- 59 42 Clausen J, Marott JL, Holtermann A, *et al.* Midlife Cardiorespiratory Fitness and the Long-Term
- 60

1
2
3 1 Risk of Mortality 46 Years of Follow-Up. *J Am Coll Cardiol* 2018;**72**:987–95.
4
5 2 doi:10.1016/j.jacc.2018.06.045
6
7 3
8
9
10 4 43 Kodama S, Saito K, Tanaka S, *et al.* Cardiorespiratory fitness as a quantitative predictor of all-
11
12 5 cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*
13
14 6 2009;**301**:2024–35. doi:10.1001/jama.2009.681
15
16 7
17
18
19 8 44 Simmons R, Griffin S, Steele R, *et al.* Increasing overall physical activity and aerobic fitness is
20
21 9 associated with improvements in metabolic risk: cohort analysis of the ProActive trial. *Diabetologia*
22
23 10 2008;**51**:787–94. doi:10.1007/s00125-008-0949-4
24
25 11
26
27
28 12 45 Department of Health. Start Active, Stay Active – A report on physical activity for health from the
29
30 13 four home countries’ Chief Medical Officers. *London: Departmet of Health* 2011.
31
32 14
33
34 15 46 Berlin JE, Storti KL, Brach JS. Using Activity Monitors to Measure Physical Activity in Free-Living
35
36 16 Conditions. *Physical Therapy* 2006;**86**:1137–45. doi:10.1093/ptj/86.8.1137
37
38
39 17
40
41 18 47 Strath SJ, Kaminsky LA, Ainsworth BE, *et al.* Guide to the assessment of physical activity: Clinical
42
43 19 and research applications: a scientific statement from the American Heart Association. *Circulation*
44
45 20 2013;**128**:2259–79. doi:10.1161/01.cir.0000435708.67487.da
46
47
48 21
49
50 22 48 Morris S, Earl K. Health Survey for England 2016 Well-being and mental health. *Health and Social*
51
52 23 *Care Information Centre* 2017.
53
54 24
55
56
57 25 49 Wade M, Mann S, Copeland RJ, *et al.* The effect of exercise referral schemes upon health and
58
59 26 wellbeing: Initial observational insights using individual patient data meta-analysis from The National
60

- 1 Referral database. doi:10.31236/osf.io/yebmr
- 2
- 3
- 4
- 5
- 6
- 7
- 8 50 Shah N, Cader M, Andrews WP, *et al.* Responsiveness of the Short Warwick Edinburgh Mental
- 9
- 10 Well-Being Scale (SWEMWBS): evaluation a clinical sample. *Health and Quality of Life Outcomes*
- 11
- 12 2018;**16**:239. doi:10.1186/s12955-018-1060-2
- 13
- 14
- 15
- 16
- 17 51 Cavill N, Roberts K, Rutter H. Standard evaluation framework for physical activity interventions.
- 18
- 19 Oxford: National Obesity Observatory 2012.
- 20
- 21
- 22
- 23
- 24 52 Craig P, Dieppe P, Macintyre S, *et al.* Developing and evaluating complex interventions: the new
- 25
- 26 Medical Research Council guidance. *BMJ* 2008;**337**:a1655. doi:10.1136/bmj.a1655
- 27
- 28
- 29
- 30 53 Rutter H, Savona N, Glonti K, *et al.* The need for a complex systems model of evidence for public
- 31
- 32 health. *Lancet* 2017;**17**:1267–9. doi:10.1016/S0140-6736
- 33
- 34
- 35
- 36
- 37 54 Beedie C, Mann S, Jimenez A, *et al.* Death by effectiveness: exercise as medicine caught in the
- 38
- 39 efficacy trap! *Brit J Sport Med* 2015;**0**:1–2. doi:10.1136/bjsports-2014-094389
- 40
- 41
- 42
- 43
- 44 55 Perera R, Heneghan C, Yudkin P. Graphical method for depicting randomised trials of complex
- 45
- 46 interventions. *BMJ* 2007;**334**:127. doi:10.1136/bmj.39045.396817.68
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Pragmatic evaluation of a co-produced physical activity referral scheme: A quasi-experimental
2 study

3 Benjamin J. R. Buckley^a, Dick H. J. Thijssen^{a,e}, Rebecca C. Murphy^a, Lee E. F. Graves^a,
4 Madeleine Cochrane^a, Fiona Gillison^b, Diane Crone^c, Philip M. Wilson^d, Greg Whyte^a and
5 Paula M. Watson^a

6
7 ^a Research Institute for Sport and Exercise Sciences, Liverpool John Moores University,
8 Liverpool, UK

9 ^b Department for Health, University of Bath, Bath, UK

10 ^c Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, UK

11 ^d Behavioural Health Sciences Research Lab, Department of Kinesiology, Brock University,
12 Ontario, Canada

13 ^e Radboud Institute for Health Sciences, Department of Physiology, Radboud University
14 Medical Center, Nijmegen, Netherlands

15
16
17 Correspondence to Dr Ben Buckley: B.J.Buckley@ljmu.ac.uk

18
19 Contributorship Statement

20 BJRB contributed to the study design, data collection, data analysis, and preparation of the final
21 document. PMW, DHJT, and RCM contributed to the study design, data analysis, and preparation of
22 the final document. MC contributed to the data collection and approved the final version. LEFG, FG,
23 DC, PW, and GW intellectually contributed to this paper and approved the final version.

Objectives. UK exercise referral schemes (ERSs) have been criticised for focusing too much on exercise prescription and not enough on sustainable physical activity (PA) behaviour change. Previously, a theoretically-grounded intervention (Co-PARS) was co-produced to support long-term PA behaviour change in individuals with health conditions. The purpose of this study was to investigate the effectiveness of Co-PARS compared to a usual care ERS and no treatment for increasing cardiorespiratory fitness.

Design. A three-arm quasi-experimental trial.

Setting. Two leisure centres providing a) Co-PARS, b) usual exercise referral care, and one no-treatment control.

Participants. 68 adults with lifestyle-related health conditions (e.g. cardiovascular, diabetes, depression) were recruited to Co-PARS, usual care, or no treatment.

Intervention. 16-weeks of physical activity behaviour change support delivered at 4, 8, 12, and 18 weeks, in addition to the usual care 12-week leisure centre access.

Outcome measures. Cardiorespiratory fitness, vascular health, PA, and mental wellbeing were measured at baseline, 12 weeks, and 6 months (PA and mental wellbeing only). Fitness centre engagement (Co-PARS and usual care) and behaviour change consultation attendance (Co-PARS) were assessed. Following an intention-to-treat approach, repeated-measures linear mixed models were used to explore intervention effects.

Results. Significant improvements in cardiorespiratory fitness ($p=.002$) and vascular health ($p=.002$) were found in Co-PARS compared to usual care and no-treatment at 12 weeks. No significant changes in PA or wellbeing at 12 weeks or 6 months were noted. Intervention engagement was higher in Co-PARS than usual care, though this was not statistically significant.

Conclusion. A co-produced PA behaviour change intervention led to promising improvements in cardiorespiratory and vascular health at 12 weeks, despite no effect for PA levels at 12 weeks or 6 months.

Trial registration: ClinicalTrials.gov: NCT03490747

Keywords: Cardiovascular Health; Self-Determination Theory; Exercise Referral; Behaviour Change Intervention; Translational Research.

1
2
3 1
4
5 2 **Strengths and limitations of the study**
6
7 3 • This study advances the literature on exercise referral effectiveness by pragmatically evaluating a
8 4 co-produced physical activity referral intervention, which was underpinned by multiple
9 5 stakeholders and behaviour change theory.
10
11 6 • The study documents the third phase of a novel and iterative approach which co-produced,
12 7 piloted, and then evaluated (this study) a physical activity referral intervention that was deemed
13 8 feasible to implement in practice.
14
15 9 • Objective and subjective measures provide insight into the potential effects for patient health.
16
17 10 • It is not possible to directly attribute intervention effects to the phased co-production approach,
18 11 although supported by the Medical Research Council.
19
20 12 • A larger sample size is needed to substantiate findings.
21
22
23
24 13
25 14 **Funding**
26
27 15 This project was supported by a PhD studentship for Benjamin Buckley from Liverpool John Moores
28 16 University. The 6-month data collection and analysis was supported by a financial grant from NHS
29 17 Liverpool Clinical Commissioning Group.
30
31
32 18 **Competing interests**
33
34 19 The authors declare that they have no competing interests.
35
36 20 **Authors' contributions**
37 21 BJRB contributed to the study design, data collection, data analysis, and preparation of the final
38 22 document. PMW, DHJT, and RCM contributed to the study design, data analysis, and preparation of
39 23 the final document. MC contributed to the data collection and approved the final version. LEFG, FG,
40 24 DC, PW, and GW intellectually contributed to this paper and approved the final version.
41
42
43 25
44 26 **Availability of data and materials**
45
46 27 The datasets used and/or analysed during the current study are available from the corresponding
47 28 author on reasonable request.
48
49 29 **Word count**
50
51 30 ~3000
52
53 31 **Ethics approval and consent to participate**
54 32 Full written consent was obtained from participants and the study was approved by NHS Research
55 33 Ethics Committee (REC: 18/NW/0039 - Project: 238547).
56
57 34 **Acknowledgements**
58
59
60

We would like to thank the participants in this study for their time, the delivery staff and centre managers for their ongoing support, and the initial development group involved in the co-production process.

INTRODUCTION

Physical inactivity is the fourth leading cause of death worldwide and costs the UK an estimated £7.4 billion annually, including £0.9 billion to the NHS alone[1]. Exercise referral schemes (ERSs) provide a promising framework to facilitate physical activity (PA) behaviour change in at-risk populations. Typically, UK ERSs consist of a referral from a healthcare professional to a 12-16-week tailored exercise programme provided by a qualified practitioner.

There is inconsistent evidence as to the effectiveness of ERSs on PA behaviour, mental well-being, quality of life, and physical health outcomes [2–4]. More recently, however, promising effects of ERSs have been demonstrated in Wales [5], Sweden [6], and Spain [7] and a systematic review identified promising effects of UK ERSs on self-reported PA and cardiovascular health markers [8]. Prior and colleagues [9] demonstrated that for every 11 participants referred to a 24-week ERS, 1 participant went on to report achieving ≥ 90 min/week of PA at 12-months. For perspective, it is estimated that 67-167 patients (categorised as $\leq 10\%$ cardiovascular disease (CVD) risk) need to receive statin treatment for 5 years to prevent one major vascular event [10]. Whilst we are not suggesting PA behaviour change is a comparable outcome to a serious clinical event, it is notable that replacing 30 minutes of TV viewing time with PA across the UK population, could reduce premature mortality by 5-15%, depending on activity intensity [11]. The majority of studies evaluating ERSs, however, have drawn on self-reported PA data and future studies employing device-based measures are needed to substantiate these observations.

Despite recent promise for the effectiveness of ERSs [7–9,12], substantial heterogeneity exists in both design and delivery [13,14], reflecting varying assumptions on how best to promote health behaviour change [15,16]. This limits potential scalability of ‘successful’ ERSs. Traditionally, ERSs have focussed on short-term exercise prescription without appropriate evidence of effectiveness or underpinning of

1
2
3 1 behaviour change theory [17]. A recent attempt to integrate behaviour change theory into an ERS [18]
4
5 2 however, showed no advantage over a standard ERS at 12 weeks or 6 months. The authors noted
6
7 3 considerable implementation challenges when training staff, such as work-related demands that may
8
9 4 have reduced the importance of the theory-based training. It is plausible that delivery staff asked to
10
11 5 implement interventions designed by academics may lack ownership and feel less
12
13 6 motivated/competent. One potential way to promote ownership and engagement might be to adopt
14
15 7 a co-production approach, as a means of co-creating value across the public sector [19–21]. Though
16
17 8 not a panacea, the involvement of practitioners, managers and service-users in co-production has
18
19 9 potential to improve intervention relevance, fidelity, and effectiveness [22].
20
21
22
23 10 Previously, a theoretically-grounded PA referral scheme (Co-PARS) was co-produced by academics,
24
25 11 policy-makers, practitioners, and service-users [23] in Liverpool, UK, with a focus on supporting
26
27 12 sustainable PA behaviour change. Liverpool is the 3rd most deprived local authority in England and
28
29 13 has the 2nd highest proportion of Lower Super Output Areas (LSOAs) in the most deprived 10%
30
31 14 nationally [24]. Interventional work with at-risk patients is therefore critical and is aligned with the
32
33 15 concept of proportionate universalism [25]. Underpinned by self-determination theory [24], the co-
34
35 16 produced intervention differed from usual ERS care in its focus on PA behaviour change (rather than
36
37 17 exercise prescription), and inclusion of frequent one-to-one consultations with exercise referral
38
39 18 practitioners (compared to usual care which included formal contact at induction only). A pilot of Co-
40
41 19 PARS [26] showed clinically meaningful improvements in cardiorespiratory fitness (CRF) and PA,
42
43 20 although as we did not include a usual care control, it was unknown whether these effects were due
44
45 21 to the fact participants were taking part in an ERS or due to the unique elements of Co-PARS.
46
47 22 Furthermore, despite having very low CRF ($<27.7 \text{ ml.kg}^{-1}.\text{min}^{-1}$) [26] we found 64% of the baseline pilot
48
49 23 sample were meeting the PA guidelines [27] of at least 150 minutes moderate-intensity PA per week
50
51 24 (measured objectively via accelerometry). This suggested CRF may be a more appropriate primary
52
53 25 outcome measure than PA for this low-fit population (whilst changing PA behaviour was the focus of
54
55 26 the intervention, a target health outcome of this behaviour change was improved CRF). The pilot also
56
57
58
59
60

1 allowed the opportunity to investigate delivery processes, and we noted several areas that required
2 refinement in preparation for a controlled trial. These refinements included, increasing the number
3 of behaviour change consultations from four to five; enhanced focus on daily PA opportunities (rather
4 than focussing on activities offered at the fitness centre); adapting staff timetables to promote
5 consistency of care and to allow participant one-to-one consultations to take place in a private room;
6 and reducing practitioner paperwork. Building on our previous pilot work, the aim of the current study
7 was to investigate the effectiveness of Co-PARS compared to a usual care ERS and a no-treatment
8 control on change in cardiorespiratory fitness (CRF) at 12 weeks and PA and wellbeing at 6 months.

9 **METHODS**

10 **Study Design**

11 A three-arm quasi-experimental trial involving: 1. Co-PARS (delivered at fitness centre A); 2. usual care
12 ERS (delivered at fitness centre B); and 3. no-treatment control. This paper reports trial outcomes
13 (CRF, vascular health, PA, mental wellbeing) measured at baseline, 12 weeks, and 6 months (PA and
14 mental wellbeing only). Additional data were collected to investigate psychosocial processes of
15 change, intervention fidelity and cost-effectiveness; due to space limitations they are not considered
16 in the present manuscript, but findings can be obtained on request from p.m.watson@ljmu.ac.uk. Full
17 written consent was obtained from participants and the study was approved by NHS Research Ethics
18 Committee (REC: 18/NW/0039 - Project: 238547) and registered on ClinicalTrials.gov (NCT03490747).

19 **Patient and Public Involvement**

20 The intervention was previously co-produced, piloted, and adapted with substantial service user input
21 [23,26].

22 **Participants and Recruitment**

23 Inclusion criteria were the same for all three conditions (Co-PARS, usual care, no-treatment).
24 Participants were eligible if aged ≥ 18 years with a health-related risk factor (e.g. hypertension,
25 hyperglycaemia, obesity) and/or health condition (e.g. diabetes, cardiovascular disease, depression)

that may be alleviated by increasing PA levels. Participants with uncontrolled health conditions, severe psychological or neurological conditions were excluded. Participants for the Co-PARS and usual care arms were recruited from fitness centre A (Co-PARS) and fitness centre B (usual care) respectively (where they had been referred for exercise by a health professional). Reception staff at both centres provided study information and gained consent to pass participant details to the researcher. Participants for the no-treatment control were recruited via posters, electronic invitations, and email communications primarily at the university site. Participants were not eligible for the no-treatment control if they were currently attending an exercise referral scheme. Interested participants for all groups were sent an information sheet and baseline data collection was arranged.

Study Arms

Intervention arm components are presented in Figure 1.

Usual care exercise referral scheme (ERS – centre B). Usual care followed a standard ERS model of 12-week subsidised access to a fitness centre (swimming, gym, group classes). Participants met an exercise referral practitioner for an initial, 1-hour induction (week 1) during which a 12-week exercise programme was provided for the participant. Any further contact with a practitioner was informal and opportunistic. This system was already in place and was considered usual care for the local area. Centre B was chosen as a comparison centre due to its similarity in referral numbers and socio-economic make-up of the local population to centre A (where Co-PARS was being delivered). For example, based on areas within Liverpool ranked from 1 (most deprived) to 30 (least deprived), usual care ERS and Co-PARS were ranked respectively: 20th and 21st (income), 20th and 21st (employment), 22nd and 24th (Education) and 10th and 11th (living environment).

Co-produced PA referral scheme (Co-PARS – centre A)

Participants received the same 12-week subsidised access to a fitness centre as usual care plus a series of one-to-one behaviour change consultations (60-minute induction followed by 30-minute consultations at weeks 4, 8, 12 and 18). A log book was provided for each participant to set action

plans, log progress and facilitate consultation discussions. Consultations were delivered by exercise referral practitioners in an autonomy supportive counselling style, drawing on the principles of self-determination theory [28]. This additional support aimed to encourage habitual opportunities to increase PA as well as activities available at the fitness centre. A full description of the theoretical underpinning and behaviour change intervention components is available elsewhere [23].

Prior to the pilot of Co-PARS [26] practitioners received training in self-determination theory-based communication strategies led by a sport and exercise psychologist (last author [PMW]), involving a workshop, one-to-one sessions and follow-up group meetings. Following the pilot, a further series of group meetings involving exercise referral practitioners and the research team were held to develop aspects of delivery that required refinement (as outlined in the introduction). Full details of the training are available from p.m.watson@ljmu.ac.uk.

No-treatment control (NTC). Participants received a lifestyle advice booklet only (offered to all study arms at baseline data collection), based on national guidance for PA, nutrition, smoking cessation and alcohol consumption.

[INSERT FIGURE 1 SOMEWHERE HERE]

Outcome measures

Primary outcome: Cardio-respiratory fitness (CRF). Maximal oxygen consumption ($\text{VO}_{2\text{max}}^2$) was estimated via the sub-maximal Astrand-Rhyming cycle ergometer protocol [29]. The protocol is a single-stage cycling test designed to elicit a steady-state heart rate over a period of ~6 minutes.

Accelerometer-derived PA. Tri-axial ActiGraph GT3x accelerometers (ActiGraph, Pensacola, FL, USA) measured PA for 7 days, which have been validated in a comparable population [30]. Raw triaxial acceleration values were converted into an omnidirectional measure of acceleration, referred to as Euclidian norm minus one [31]. Minimum wear time was 10 hours per day and 3 days per week including one weekend day [32]. The R package GGIR [31] facilitated extraction of user-defined

1
2
3 1 acceleration thresholds: 5.9 to 69.1 mg for light-intensity PA [33], 69.1 to 258.7 mg as moderate and
4
5 2 >258.7 mg as vigorous-intensity PA [34].
6
7
8 3 *Vascular health.* Our previous work has demonstrated carotid artery reactivity (CAR) may be a
9
10 4 promising outcome variable to assess in PA interventions for at-risk populations [35]. Further,
11
12 5 endothelial function may provide prognostic value beyond that of traditional risk factors [36] with an
13
14 6 increase of 1% in brachial artery flow-mediated dilation (FMD) associated with a 12-15% lower risk of
15
16 7 CV events [33,34]. FMD and CAR were measured using ultrasound techniques [35]. Both techniques
17
18 8 measure vascular endothelial function and have independently predicted future risk of cardiovascular
19
20 9 events in humans [36,37]. Blood pressure was measured in the supine position using an automated
21
22 10 blood pressure device (Omron Healthcare UK Limited, Milton Keynes, UK).
23
24
25
26 11 *Anthropometric measures.* Since obesity is a critical risk factor for poor health and cardiovascular
27
28 12 disease, anthropometric variables were measured to investigate potential intervention effects on
29
30 13 body mass. Waist-to-height ratio is a stronger predictor of early health risk than Body Mass Index
31
32 14 (BMI) alone [38], therefore we collected both BMI (mass in kg / stature in m²) and waist-to-height
33
34 15 ratio (waist circumference / stature).
35
36
37
38 16 *Mental wellbeing.* As PA is known to enhance mental wellbeing [39] and clinical populations are more
39
40 17 susceptible to mental ill-health [40], it was important to identify whether Co-PARS led to any changes
41
42 18 in mental health (positive or negative). Mental wellbeing was measured using the 14-item Warwick-
43
44 19 Edinburgh Mental Well-being Scale (WEMWBS; [41], which asks participants to rate their
45
46 20 psychological wellbeing (e.g. “I’ve been feeling cheerful”) over the previous 2 weeks (measured on a
47
48 21 likert scale of 1 (none of the time) to 5 (all of the time)).
49
50
51
52 22 *Fitness centre engagement (Co-PARS and usual care only).* The number of occasions participants
53
54 23 attended the fitness centre between baseline and 12 weeks (weekly attendance) and 12 weeks to 6
55
56 24 months (monthly attendance) was obtained from computerised attendance records. When
57
58 25 measuring intervention engagement it was deemed inappropriate to calculate the mean number of
59
60

sessions per week, since this could exaggerate the engagement of individuals who attended with high frequency in the early weeks then dropped out (when compared with individuals who attended moderately but consistently for the full 12 weeks). Therefore a formula was used to calculate a percentage for '12-week engagement' (based on the recommended bi-weekly attendance):

$$\left(\frac{((n1*0.5) + (n2) + (n3*1.2))}{12} \right) * 100$$

n1 = number of weeks in which participant attends once only
n2 = number of weeks in which participant attends twice
n3 = number of weeks in which participant attends three or more times

This formula took into account both *frequency* and *consistency* of attendance to yield a percentage score that ranged from 0% (no attendance) to 120% (attendance of three or more times per week for the whole 12 weeks).

Monthly attendance post-12 weeks was calculated as a mean attendance across months 4 to 6, therefore did not take consistency of attendance into account.

Behaviour change consultation attendance (Co-PARS only). The number of consultations offered and attended were measured by exercise referral practitioners at induction, 4, 8, 12, and 18 weeks.

Sample size

Sample size was determined to detect a meaningful difference in CRF at 12 weeks based on our pilot results [26]. To detect a difference of 2 ml.kg⁻¹min⁻¹ between Co-PARS and usual care, 42 participants were required per arm (f= .25, p= .05, power = .80). To detect a difference of 3.2 ml.kg⁻¹min⁻¹ between the intervention arms and the no-treatment control, 17 participants were required for the no-treatment control (f= .5, p= .05, power = .80). Thus, a total sample of 101 participants were required.

Statistical analyses

An intention-to-treat approach was used assuming no change in non-respondents (last observation carried forward) to produce a conservative estimate of intervention effects. Delta changes (Δ) from

1
2
3 1 pre- to post-intervention were calculated for each group and entered as the dependent variable in
4
5 2 repeated measures linear mixed model analyses. A random intercept model was used with fixed
6
7 3 effects for study arm (Co-PARS, usual care ERS, no-treatment control) and time (baseline-to-week-12
8
9 4 change, week-12-to-6-month change, and baseline-to-6-month change) and participants included as
10
11 5 random effects. Least squared difference (LSD) was used for post hoc testing. Testing for baseline
12
13 6 differences to identify covariates was avoided, as this method has been demonstrated to inflate bias,
14
15 7 instead pre-intervention was entered into the model as a covariate. Furthermore, all linear mixed
16
17 8 model analyses were repeated with age and employment as covariates as a comparison to the results
18
19 9 presented in this study (with baseline score as a covariate) due to their known prognostic value. Using
20
21 10 age and employment as covariates resulted in no change in inferences presented in this study. One-
22
23 11 way ANOVAs were used to compare baseline values between intervention arms. Fitness centre
24
25 12 engagement was determined as described above. Behaviour change consultation attendance is
26
27 13 presented descriptively. For non-normally distributed data, median and interquartile range is
28
29 14 presented and within group median change was calculated via Wilcoxon signed-rank tests.
30
31
32
33
34

35 15 **RESULTS**

36
37
38 16 *Participants.* 68 participants provided baseline data, 56 of whom provided 12-week data, and 58 of
39
40 17 whom provided 6-month data (figure 2).
41
42

43 18 **Baseline characteristics (table 1).** No significant differences were noted between arms for age, sex,
44
45 19 ethnicity, BMI, referral reason, or accelerometer-derived PA levels ($p>.05$). Full-time employment
46
47 20 status ($p=.001$) and CRF ($p=.015$) were significantly higher in the control compared to usual care and
48
49 21 Co-PARS. Smoking status was significantly higher in usual care compared to Co-PARS and control
50
51 22 ($p=.010$). Mental wellbeing was significantly lower in Co-PARS compared to control ($p=.023$).
52
53
54
55

56 23
57
58 24 [INSERT FIGURE 2 SOMEWHERE HERE]
59
60

Table 1. Baseline characteristics presented as Mean \pm SD or % (n) of group.

	Co-produced PA referral (n=33)	Usual care ERS (n=19)	No-treatment control (n=16)	Between arm <i>p</i> -value
Age (years)	57 \pm 12	53 \pm 16	48 \pm 15	<i>p</i> =.319
Female (% of sample)	58 (19)	47 (9)	56 (9)	<i>p</i> =.774
White British (% of sample)	82 (27)	95 (18)	75 (12)	<i>p</i> =.132
Full-time employment (% of sample)	18 (6)	26 (5)	62 (10)	<i>p</i> =.001
Never smoked (% of sample)	73 (24)	37 (7)	81 (13)	<i>p</i> =.002
Body mass index (kg/m ²)	31 \pm 7	33 \pm 6	29 \pm 6	<i>p</i> =.226
Systolic blood pressure (mmHg)	131 \pm 11	138 \pm 18	123 \pm 12	<i>p</i> =.010
Primary referral reason / health concern (control)				<i>p</i> =.132
Cardiometabolic (% of sample)	67 (22)	43 (8)	62 (10)	-
Cancer (% of sample)	6 (2)	5 (1)	6 (1)	-
Mental Health (% of sample)	18 (6)	26 (5)	19 (3)	-
Musculoskeletal (% of sample)	9 (3)	26 (5)	13 (2)	-
Comorbidity (% of sample)	85 (28)	100 (19)	81 (13)	<i>p</i> =.166
Meeting the PA guidelines (% of sample)*	73 (22)	71 (10)	93 (13)	<i>p</i> =.223

P-values represent between arm baseline effects. There was no between arm effect for referral reason, thus no between arm *p*-values are provided for referral reason sub groups.

*Chief Medical Officers' 2019 physical activity guidelines: 150 minutes of moderate-intensity physical activity per week.

Baseline-to-12-Week effects

Raw outcome values are presented for baseline, week 12, and 6 months in Table 2. There was a significant effect for study arm in baseline-to-12-week change in CRF (*p*=.002). Post hoc testing

1
2
3 1 revealed a significantly higher CRF change in Co-PARS (2.4) compared to the ERS (0.3; $p=.021$) and
4
5 2 control (-0.6; $p=.001$), but no difference between the ERS and control ($p=.314$). A significant effect for
6
7 3 study arm was found in change in FMD% ($p=.002$), with FMD% change significantly higher in Co-PARS
8
9 4 (2.4) compared to control (-1.1; $p=.001$) but not the ERS (0.8; $p=.099$). The change in FMD% was not
10
11 5 significantly different between the ERS and control ($p=.71$). No statistically significant study arm
12
13 6 effects were noted for changes in CAR%, blood pressure, resting heart rate, anthropometric measures,
14
15 7 PA or WEMWBS at 12 weeks ($p>.05$).

18
19
20 8 **Baseline-to-6-month effects**

21
22 9 No statistically significant study arm effects were noted for change in WEMWBS or PA at 6 months
23
24 10 ($p>.05$).

25
26
27
28 11 **Fitness centre engagement (Co-PARS and usual care ERS) and consultation attendance (Co-PARS**
29
30 12 **only).**

31
32
33 13 Table 3 reports the participant fitness centre engagement data for the Co-PARS and usual care ERS.
34
35 14 Although not statistically significant, Co-PARS engagement was 9% higher, participants attended the
36
37 15 fitness centre on average 3 times more per month, and 23% more participants were attending the
38
39 16 fitness centre beyond 6-months follow-up compared to usual care. Co-PARS behaviour change
40
41 17 consultation attendance is reported in Table 4.

42
43
44
45
46 18

Table 2. Cardiometabolic health outcomes and PA levels at baseline, 12 weeks, 6 months, and between arm baseline-to 12-week or 6-month effect. All variables are presented as Mean \pm SD.

	Co-PARS			Usual Care ERS			No-Treatment Control			Between arm effect <i>p</i> -value ^(a)
	Baseline	Week 12	6 Month	Baseline	Week 12	6 Month	Baseline	Week 12	6 Month	
Fitness (n=56)										
CRF _{ml.kg.⁻¹min⁻¹}	22.2±7	24.6±7	-	23.3±6.6	23.6±7	-	29.6±9.2	28.9±8.7	-	<i>p</i> =.002
Physical Activity										
GT3x (n= 61) _{Mins.day}										
Light intensity	90±52	98±64	107±75	98±36	93±31	158±145	90±37	101±33	86±40	<i>p</i> =.332
Moderate intensity	44±32	42±29	42±33	43±28	43±30	55±55	60±31	65±24	54±21	<i>p</i> =.260
Vigorous intensity	1±3	1±2	1±2	1±2	1±1	1±2	2±4	2±3	3±8	<i>p</i> =.108
Vascular Ultrasound (n=64)										
CAR%	1.7±2.7	2.8±2.2	-	2.7±1.8	3.9±2.8	-	2.5±2.7	1.7±2.7	-	<i>p</i> =.073
CAR Baseline _{cm}	0.69±0.07	0.69±0.06	-	0.69±0.08	0.7±0.09	-	0.65±0.07	0.64±0.06	-	<i>p</i> =.130
FMD%	4.4±2.3	6.8±2.7	-	4.2±2	5±2.1	-	6.2±2.1	5.2±2.8	-	<i>p</i> =.002
FMD Baseline _{cm}	0.39±0.07	0.38±0.06	-	0.39±0.09	0.41 0.08	-	0.38±0.08	0.37±0.06	-	<i>p</i> =.728
Cardiometabolic (n=68)										
BMI _{kg.m²}	31±7	30±7	-	33±6	32±6	-	29±6	29±6	-	<i>p</i> =.323
WHR	62±9	61±10	-	64±8	63±8	-	56±9	56±9	-	<i>p</i> =.261
SBP _{mmHg}	131±11	127±12	-	138±18	132±15	-	123±12	118±13	-	<i>p</i> =.937
DBP _{mmHg}	73±7	71±8	-	73±9	71±11	-	72±11	68±10	-	<i>p</i> =.584
RHR _{bpm}	70±10	65±10	-	70±12	68±11	-	66±12	63±9	-	<i>p</i> =.540
Mental Wellbeing (n=68)										
WEMWBS	46±9	51±10	48±10	49±10	52±11	50±13	53±9	56±9	53±10	<i>p</i> =.796

Co-PARS, Co-produced PA referral scheme; ERS, Exercise referral scheme; CRF, Cardiorespiratory Fitness; GT3x, Accelerometer; CAR, Carotid artery reactivity; FMD, Flow-mediated dilation; BMI, Body Mass Index; WHR, Waist-to-Height ratio; SBP, Systolic blood pressure; DBP, Diastolic blood pressure; RHR, Resting heart rate, WEMWBS, Warwick-Edinburgh Mental Wellbeing Scale

^a F-statistic for between arm baseline-to-6-month change or baseline-to-week 12 change if variable not collected at 6 months.

Missing data was due to inability to complete the CRF test (n=12), inability to complete the vascular ultrasound protocols (n=4), and insufficient accelerometer wear time or non-return (n=7).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 3. Fitness centre engagement.

	Co-PARS (n=33)	Usual Care (n=19)	Between centre difference
% Engagement ^a (Mean ± SD)	42±29	33±27	p=.267
Number of fitness centre visits (per person per month) week 12 to 6 months (Med, IQR)	3(0-14)	0 (0-1)	p=.072
% of baseline sample who attended fitness centre at least once beyond 6 months (% of sample, n)	39 (13)	16 (3)	p=.101

^aBased on the formula $((n1*0.5)+(n2)+(n3*1.2))/12 * 100$; n1=number of weeks in which participant attends once only; n2=number of weeks in which participant attends twice; n3=number of weeks in which participant attends three or more times. ^a**Engagement**; based on a recommended attendance of twice weekly, a formula was used to calculate a percentage for “12-week engagement”, which took into account both frequency and consistency of attendance (see methods).

Table 4. Co-PARS behaviour change consultation attendance (based on baseline sample of 33 participants).

Consultation	% Booked (n)	% Attended (n)
Induction	91(30)	93(28)
Week 4	82(27)	78(21)
Week 8	67(22)	91(20)
Week 12	64(21)	81(17)
Week 18	55(18)	50(9)

DISCUSSION

This was the first study to investigate the effectiveness of a theoretically-grounded, co-produced PA referral scheme (Co-PARS) compared to a usual care ERS and no treatment. Despite challenges in recruitment that meant the study was statistically underpowered, the findings demonstrated significant and clinically meaningful improvements in CRF and vascular health in Co-PARS compared to the usual care and no treatment. No statistically significant effects were noted for accelerometer-derived PA levels or mental wellbeing at 12-weeks or 6-months.

The effect of usual care ERSs compared to theoretically-grounded interventions on CRF has not been previously explored. We observed a significant increase in CRF in Co-PARS compared to usual care and a no-treatment control. According to values reported by Clausen *et al.* [42] both Co-PARS ($22 \text{ ml.kg}^{-1}\text{min}^{-1}$) and usual care ($23 \text{ ml.kg}^{-1}\text{min}^{-1}$) participants were below the lower limit of 'healthy' ($27.7 \text{ ml.kg}^{-1}\text{min}^{-1}$) for baseline CRF [43]. As low CRF is associated with a substantially elevated risk of all-cause mortality [43], the magnitude of change demonstrated in Co-PARS ($2.4 \text{ ml.kg}^{-1}\text{min}^{-1}$) may be clinically meaningful. For example, in at-risk populations, relatively small magnitudes ($\leq 1 \text{ ml.kg}^{-1}\text{min}^{-1}$) have been shown to significantly reduce clustered cardiometabolic risk [44]. Thus, Co-PARS was effective at improving CRF in individuals with low CRF by a clinically meaningful amount.

Promising improvements in vascular health were also noted in the Co-PARS group, with brachial artery FMD significantly improved compared to usual care and control arms. Although CAR was not statistically different between arms, both Co-PARS and usual care demonstrated a potentially meaningful within-arm improvement compared with no treatment, which exhibited a deterioration in vascular health. Such improvements in vascular measures may have prognostic implications. For example, a 1% increase in FMD has been suggested to reduce the future risk of CVD events by 13% [36].

Despite low baseline CRF, a substantial percentage of Co-PARS (73%) and usual care (71%) participants were meeting the Department of Health [45] guidelines of 150 minutes of moderate-intensity PA per

1
2
3 1 week. We observed a similar finding in our pilot [26] and subsequently raised the question as to the
4
5 2 use of PA guidelines to assess eligibility for ERSs (NICE, 2014), as it appears from our data that
6
7 3 individuals classified as “physically active” can still be very unfit and therefore can benefit from ERSs
8
9
10 4 in terms of improved fitness and cardiometabolic health. A further discrepancy was noted in the lack
11
12 5 of change in PA levels in Co-PARS, despite improved CRF. It is possible measurement issues
13
14 6 contributed to this discrepancy. Accelerometers can measure certain types of PA such as walking,
15
16 7 running, and stair climbing [46]. They may not, however, sufficiently identify activities typical of an
17
18 8 ERS delivered within a fitness centre environment (e.g. cycling, resistance training, circuits,
19
20 9 swimming). Given Co-PARS had higher (albeit non-significant) fitness centre engagement compared
21
22
23 10 to usual care, it is possible PA changes occurred that were not detected by the accelerometry data.
24
25 11 Consideration therefore needs to be given to the appropriateness of accelerometers to measure PA
26
27 12 in ERSs. Alternative methods such as heart-rate monitors combined with self-report data may be
28
29 13 worthy of consideration, although further work would be required to develop standardized data
30
31 14 collection and analysis protocols (taking into account the limitations of each of these methods if used
32
33 15 in isolation [47]). Researchers are therefore urged to consider CRF as a primary outcome in ERSs until
34
35 16 appropriate alternative methods of measuring PA behaviour are developed. Ultimately, it is not clear
36
37 17 why the increase in fitness occurred without a corresponding change in PA and further research is
38
39 18 required to elucidate the relationship between PA and fitness in this population.
40
41
42
43
44 19 In addition to physiological health outcomes, we found baseline mental wellbeing to be below the
45
46 20 national average (score of 50) in both Co-PARS (46) and usual care (49), but not the control (53) [48].
47
48 21 Despite no significant between-group effect for mental wellbeing, within-group changes at 12 weeks
49
50 22 were deemed clinically meaningful for Co-PARS (5) and usual care (3) but not in the no treatment
51
52 23 control. It is notable that the post-intervention magnitude of change observed in mental wellbeing for
53
54 24 Co-PARS (5) was larger than that observed in a meta-analysis encompassing >23,000 participants
55
56 25 across 13 different ERSs (3), which were comparable in nature to the usual care ERS in this study [49].
57
58
59
60

1 From the 6-month data it appeared the scheme was not effective at promoting *sustained* PA behaviour
2 change or mental wellbeing improvements. It must be noted, however, that the wellbeing levels were
3 still higher than baseline and even small magnitudes of change (1-3) may be meaningful in clinical
4 populations [50]. As discussed earlier, it may be that measuring PA using the methods described in
5 this study prevented the identification of activities typical of a fitness centre environment. This notion
6 is supported by the post-week-12 attendance data, which highlighted Co-PARS participants were
7 regularly attending the fitness centre whereas the usual care participants were not. Challenges of
8 maintaining sustained health outcomes post-ERSs have been highlighted elsewhere [3]. And whilst a
9 recent systematic review reported longer length schemes (>20 weeks) may be more effective than
10 shorter schemes [8], the four long ERSs (20-26 weeks) collected pre-post data only. Thus we do not
11 know if longer length ERSs result in enhanced health outcomes *post intervention* compared with
12 shorter schemes. To determine if longer length schemes are indeed more effective, longer-term
13 follow-up data collection is required, ideally at 6 and 12 months post intervention [51].

14 Through a phased approach we have assessed the effectiveness of Co-PARS resulting from several
15 years of co-production. Whilst the effects of co-production are difficult to isolate, a comparison of
16 results at different stages of intervention refinement suggests the phased development approach had
17 some positive effects. Unpublished engagement data from centre A in 2014-2015 (when the centre
18 was running a usual care ERS) shows that engagement improved after the introduction of Co-PARS
19 (42% vs 28% in 2014-2015), whereas engagement reduced in the usual care centre over the same
20 period (32% vs 37% in 2014-2015). Furthermore, consultation attendance for Co-PARS in the current
21 study was substantially higher than in our previous pilot (54% attended induction plus ≥ 3 behaviour
22 change consultations, vs 9% in the pilot [26]), which may have been a reflection of refinements made
23 to the intervention after the pilot (e.g. improved focus on holistic PA, improved monitoring
24 procedures, improved continuity of instructors). These improvements in engagement highlight the
25 importance of allowing time for complex interventions to develop [52], and are particularly promising
26 given the effectiveness of ERSs are highly dependent on participant adherence [5,21]. Furthermore,

1
2
3 1 this study has demonstrated how investing in the “bottom-up” development of an intervention can
4
5 2 lead to an effective and sustainable model. We therefore support the arguments of Rutter and
6
7 3 colleagues [53] in that a shift in thinking is needed, instead of asking whether an intervention works
8
9 4 to fix a problem, researchers should aim to identify if and how it contributes to reshaping a system in
10
11 5 a favourable way. As such, we propose the co-production and implementation process may be as
12
13 6 important as the scheme content itself.
14
15

16
17 7 **Methodological considerations**
18
19

20 8 This is the first known study to investigate the effectiveness of a co-produced PA referral scheme (Co-
21
22 9 PARS) in comparison to usual care and a no-treatment control. Our novel approach addresses an
23
24 10 important gap in the sport and exercise medicine literature [54], in that we employed rigorous
25
26 11 laboratory-based instruments to measure health outcomes that can be achieved through an
27
28 12 ecologically valid, “real-world” intervention. We observed a very high retention at 6-month follow up,
29
30 13 which may be due in part to the fact many of the participants were retired (and therefore may have
31
32 14 more available time). It is possible also that the high retention was facilitated by the co-production
33
34 15 process, which involved ongoing relationships between the research and delivery teams (and
35
36 16 therefore helped with the logistics of returning accelerometers for the co-PARS and usual care
37
38 17 groups). Whilst this paper highlights many strengths of co-production, we do not wish to present co-
39
40 18 production as a panacea [19] and it is important potential challenges and costs are considered prior
41
42 19 to undertaking such an approach [21,22].
43
44
45
46
47

48 20 We must acknowledge some limitations of the study. Whilst there is a need for high-quality RCTs of
49
50 21 theoretically informed approaches to PA behaviour change [3], several pragmatic reasons meant an
51
52 22 RCT approach was not appropriate for the present study. Firstly, it was important participants could
53
54 23 choose the most convenient fitness centre. Secondly, it was important we continued work with the
55
56 24 same fitness centre and staff (following co-production [23] and pilot [26] phases) in order to develop
57
58 25 the intervention to the point where it was deemed to have a worthwhile effect [52]. A pragmatic
59
60

research approach was therefore deemed most appropriate to evaluate Co-PARS with high ecological validity. Pragmatic constraints (e.g. fitness centre refurbishments, staff illness) did however mean the required sample size was not achieved, thus inferences of effectiveness need to be taken with caution. This is particularly true for the PA data, where the relatively high variability (compared with CRF) may have contributed to the lack of change observed in PA in this study. It is recommended future work considers pragmatic risks and contingencies when planning recruitment and plans sufficient time to cope with recruitment delays. For pragmatic reasons, not all outcomes were collected at 6-months follow-up and further research is needed to collect long-term, objective health data following PA referral schemes. Finally, it must be noted that while the trial registration appears to be retrospective (April 6th 2018), the initial submission was several months prior to this (January 11th 2018). Final sign-off was delayed due to capacity issues within the research team.

CONCLUSION

A co-produced, theoretically-grounded PA referral scheme (Co-PARS) led to improved CRF and vascular health in at-risk individuals when compared to usual care and no treatment. In addition, clinically meaningful improvements in vascular health and mental wellbeing were observed at 12-weeks in both Co-PARS and usual care, but not the no treatment control group. Of note, PA remained unchanged at 12-weeks and 6-months follow-up. Adopting a phased approach has enabled multi-stakeholder input and ongoing intervention refinement, resulting in an intervention that showed promising effects on engagement and clinically meaningful improvements to participant health.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends

- Figure 1.** ‘PaT Plot’ describing intervention arm components.[55]
- Figure 2.** Participant flow diagram within the three study arms (March 2018-January 2019).
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

For peer review only

References

- 1 Public Health England. Physical activity: applying All Our Health. 2019.
- Dugdill L, Graham RC, McNair F. Exercise referral: the public health panacea for physical activity promotion? A critical perspective of exercise referral schemes; their development and evaluation. *Ergonomics* 2005;**48**:1390–410. doi:10.1080/00140130500101544
- Pavey T, Taylor A, Fox K, *et al.* Effect of exercise referral schemes in primary care on physical activity and improving health outcomes: systematic review and meta-analysis. *Bmj* 2011;**343**:d6462. doi:10.1136/bmj.d6462
- Pavey T, Anokye N, Taylor A, *et al.* The clinical effectiveness and cost-effectiveness of exercise referral schemes: a systematic review and economic evaluation. *Health technology assessment (Winchester, England)* 2011;**15**:i–xii, 1–254. doi:10.3310/hta15440
- Murphy SM, Edwards RT, Williams N, *et al.* An evaluation of the effectiveness and cost effectiveness of the National Exercise Referral Scheme in Wales, UK: a randomised controlled trial of a public health policy initiative. *Journal of epidemiology and community health* 2012;**66**:745–53. doi:10.1136/jech-2011-200689
- Onerup A, Arvidsson D, Blomqvist Å, *et al.* Physical activity on prescription in accordance with the Swedish model increases physical activity: a systematic review. *Br J Sports Med* 2018;;bjsports-2018-099598. doi:10.1136/bjsports-2018-099598
- Martín-Borràs C, Giné-Garriga M, Puig-Ribera A, *et al.* A new model of exercise referral scheme in primary care: is the effect on adherence to physical activity sustainable in the long term? A 15-

1
2
3 1 month randomised controlled trial. *BMJ Open* 2018;**8**:e017211. doi:10.1136/bmjopen-2017-017211
4
5 2
6
7 3 8 Rowley N, Mann S, Steele J, *et al.* The effects of exercise referral schemes in the United Kingdom in
8
9 those with cardiovascular, mental health, and musculoskeletal disorders: a preliminary systematic
10 4
11 review. *BMC Public Health* 2018;**18**:949. doi:10.1186/s12889-018-5868-9
12 5
13
14 6
15
16 7 9 Prior F, Coffey M, Robins A, *et al.* Long-Term Health Outcomes Associated With an Exercise
17
18 Referral Scheme: An Observational Longitudinal Follow-Up Study. *Journal of Physical Activity and*
19 8
20 *Health* 2019;**1**:1–6. doi:10.1123/jpah.2018-0442
21 9
22
23 10
24
25 11 10 Taylor F, Huffman MD, Macedo A, *et al.* Statins for the primary prevention of cardiovascular
26
27 disease. *Cochrane Db Syst Rev* 2013;**1**:CD004816. doi:10.1002/14651858.cd004816.pub5
28 12
29
30 13
31
32 14 11 Wijndaele K, Sharp SJ, Wareham NJ, *et al.* Mortality Risk Reductions from Substituting Screen
33
34 Time by Discretionary Activities. *Med Sci Sport Exer* 2017;**49**:1111–9.
35 15
36 doi:10.1249/mss.0000000000001206
37 16
38
39 17
40
41 18 12 Craike M, Wiesner G, Enticott J, *et al.* Equity of a government subsidised exercise referral scheme:
42
43 A population study. *Social Science & Medicine* Published Online First: 2018.
44 19
45 doi:10.1016/j.socscimed.2018.09.023
46 20
47
48 21
49
50 22 13 Craig A, Dinan S, Smith A, *et al.* Exercise Referral Systems: A National Quality Assurance
51
52 Framework. *Department of Health: London* Published Online First: 2001.discovery.ucl.ac.uk
53 23
54
55 24
56
57 25 14 Pavey T, Taylor A, Hillsdon M, *et al.* Levels and predictors of exercise referral scheme uptake and
58
59 adherence: a systematic review. *J Epidemiol Commun H* 2012;**66**:737–44. doi:10.1136/jech-2011-
60

- 1 200354
- 2
- 3
- 4
- 5
- 6
- 7
- 8 15 Littlecott HJ, Moore GF, Moore L, *et al.* Psychosocial mediators of change in physical activity in
- 9 the Welsh national exercise referral scheme: secondary analysis of a randomised controlled trial.
- 10
- 11
- 12 5 *International Journal of Behavioral Nutrition and Physical Activity* 2014;**11**:1–11.
- 13
- 14 6 doi:10.1186/s12966-014-0109-9
- 15
- 16
- 17 7
- 18
- 19 8 16 Hanson CL, Oliver EJ, Dodd-Reynolds CJ, *et al.* How do participant experiences and characteristics
- 20 influence engagement in exercise referral? A qualitative longitudinal study of a scheme in
- 21
- 22
- 23 10 Northumberland, UK. *Bmj Open* 2019;**9**:e024370. doi:10.1136/bmjopen-2018-024370
- 24
- 25
- 26 11
- 27
- 28 12 17 Sowden S, Raine R. Running along parallel lines: how political reality impedes the evaluation of
- 29 public health interventions. A case study of exercise referral schemes in England. *Journal of*
- 30
- 31
- 32 14 *epidemiology and community health* 2008;**62**:835–841. doi:10.1136/jech.2007.069781
- 33
- 34
- 35 15
- 36
- 37 16 18 Duda JL, Williams GC, Ntoumanis N, *et al.* Effects of a standard provision versus an autonomy
- 38 supportive exercise referral programme on physical activity, quality of life and well-being indicators:
- 39
- 40
- 41 18 a cluster randomised controlled trial. *The international journal of behavioral nutrition and physical*
- 42
- 43 19 *activity* 2014;**11**:10. doi:10.1186/1479-5868-11-10
- 44
- 45
- 46 20
- 47
- 48 21 19 Ostrom E. Crossing the great divide: Coproduction, synergy, and development. *World*
- 49
- 50 22 *Development* 1996;**24**:1073–87. doi:10.1016/0305-750x(96)00023-x
- 51
- 52
- 53 23
- 54
- 55 24 20 Clarke D, Jones F, Harris R, *et al.* What outcomes are associated with developing and
- 56
- 57 25 implementing co-produced interventions in acute healthcare settings? A rapid evidence synthesis.
- 58
- 59 26 *BMJ open* 2017;**7**:e014650. doi:10.1136/bmjopen-2016-014650
- 60

1
2
3 1
4
5 2 21 Farrance C, Tsofliou F, Clark C. Adherence to community based group exercise interventions for
6
7
8 3 older people: A mixed-methods systematic review. *Preventive medicine* 2016;**87**:155–66.
9
10 4 doi:10.1016/j.ypmed.2016.02.037
11
12 5
13
14 6 22 Rycroft-Malone J, Burton CR, Bucknall T, *et al.* Collaboration and Co-Production of Knowledge in
15
16 7 Healthcare: Opportunities and Challenges. *International journal of health policy and management*
17
18 8 2016;**5**:221–3. doi:10.15171/ijhpm.2016.08
19
20
21 9
22
23 10 23 Buckley B, Thijssen D, Murphy R, *et al.* Making a move in exercise referral: co-development of a
24
25 11 physical activity referral scheme. *Journal of Public Health* Published Online First: 2018.
26
27 12 doi:10.1093/pubmed/fdy072
28
29
30 13
31
32 14 24 The English indices of deprivation 2019. 2019.
33
34 15
35
36 16 25 Carey G, Crammond B, Leeuw E. Towards health equity: a framework for the application of
37
38 17 proportionate universalism. *Int J Equity Health* 2015;**14**:81. doi:10.1186/s12939-015-0207-6
39
40
41 18
42
43 19 26 Buckley BJ, Thijssen DH, Murphy RC, *et al.* Preliminary effects and acceptability of a co-produced
44
45 20 physical activity referral intervention. *Health Educ J* 2019;:001789691985332.
46
47 21 doi:10.1177/0017896919853322
48
49
50 22
51
52 23 27 Department of Health & Social Care. UK Chief Medical Officers’ Physical Activity Guidelines. 2019.
53
54 24
55
56 25 28 Ryan R, Deci E. Self-determination theory and the facilitation of intrinsic motivation, social
57
58 26 development, and well-being. *The American psychologist* 2000;**55**:68–78.
59
60

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

29 Astrand I. Aerobic work capacity in men and women with special reference to age. *Acta physiologica Scandinavica Supplementum* 1960;**49**:1–92.

30 Kelly LA, McMillan DG, Anderson A, *et al.* Validity of actigraphs uniaxial and triaxial accelerometers for assessment of physical activity in adults in laboratory conditions. *BMC Medical Physics* 2013;**13**:1–7. doi:10.1186/1756-6649-13-5

31 Hees VT, Gorzelniak L, Leon E, *et al.* Separating movement and gravity components in an acceleration signal and implications for the assessment of human daily physical activity. *PloS one* 2013;**8**:e61691. doi:10.1371/journal.pone.0061691

32 Matthews CE, Hagströmer M, Pober DM, *et al.* Best Practices for Using Physical Activity Monitors in Population-Based Research. *Medicine & Science in Sports & Exercise* 2012;**44**:S68. doi:10.1249/MSS.0b013e3182399e5b

33 Bakrania K, Yates T, Rowlands AV, *et al.* Intensity Thresholds on Raw Acceleration Data: Euclidean Norm Minus One (ENMO) and Mean Amplitude Deviation (MAD) Approaches. *PLOS ONE* 2016;**11**:e0164045. doi:10.1371/journal.pone.0164045

34 Hilded M, Hees Vt, Hansen B, *et al.* Age Group Comparability of Raw Accelerometer Output from Wrist- and Hip-Worn Monitors. *Medicine & Science in Sports & Exercise* 2014;**46**:1816. doi:10.1249/mss.0000000000000289

35 Buckley B, Watson PM, Murphy RC, *et al.* Carotid artery function is restored in subjects with elevated cardiovascular disease risk following a 12-week physical activity intervention. *Canadian*

1
2
3 1 *Journal of Cardiology* Published Online First: 2018. doi:10.1016/j.cjca.2018.10.015
4
5 2
6
7 3 36 Inaba Y, Chen JA, Bergmann SR. Prediction of future cardiovascular outcomes by flow-mediated
8
9
10 4 vasodilatation of brachial artery: a meta-analysis. *The international journal of cardiovascular imaging*
11
12 5 2010;**26**:631–40. doi:10.1007/s10554-010-9616-1
13
14 6
15
16 7 37 van Mil A, Pouwels S, Wilbrink J, *et al.* Carotid Artery Reactivity Predicts Events in Peripheral
17
18 8 Arterial Disease Patients. *Annals of Surgery* 2017. doi:10.1097/SLA.0000000000002558
19
20 9
21
22
23 10 38 Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist
24
25 11 circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis.
26
27 12 *Obesity reviews : an official journal of the International Association for the Study of Obesity*
28
29 13 2012;**13**:275–86. doi:10.1111/j.1467-789X.2011.00952.x
30
31 14
32
33
34 15 39 Paluska SA, Schwenk TL. Physical Activity and Mental Health. *Sports Med* 2000;**29**:167–80.
35
36 16 doi:10.2165/00007256-200029030-00003
37
38
39 17
40
41 18 40 Barnett K, Mercer SW, Norbury M, *et al.* Epidemiology of multimorbidity and implications for
42
43 19 health care, research, and medical education: a cross-sectional study. *Lancet* 2012;**380**:37–43.
44
45 20 doi:10.1016/s0140-6736(12)60240-2
46
47 21
48
49
50 22 41 Tennant R, Hiller L, Fishwick R, *et al.* The Warwick-Edinburgh Mental Well-being Scale
51
52 23 (WEMWBS): development and UK validation. *Health Qual Life Out* 2007;**5**:1–13. doi:10.1186/1477-
53
54 24 7525-5-63
55
56 25
57
58
59 26 42 Clausen J, Marott JL, Holtermann A, *et al.* Midlife Cardiorespiratory Fitness and the Long-Term
60

- 1 Risk of Mortality 46 Years of Follow-Up. *J Am Coll Cardiol* 2018;**72**:987–95.
- 2 doi:10.1016/j.jacc.2018.06.045
- 3
- 4 43 Kodama S, Saito K, Tanaka S, *et al*. Cardiorespiratory fitness as a quantitative predictor of all-
- 5 cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*
- 6 2009;**301**:2024–35. doi:10.1001/jama.2009.681
- 7
- 8 44 Simmons R, Griffin S, Steele R, *et al*. Increasing overall physical activity and aerobic fitness is
- 9 associated with improvements in metabolic risk: cohort analysis of the ProActive trial. *Diabetologia*
- 10 2008;**51**:787–94. doi:10.1007/s00125-008-0949-4
- 11
- 12 45 Department of Health. Start Active, Stay Active – A report on physical activity for health from the
- 13 four home countries' Chief Medical Officers. *London: Departmet of Health* 2011.
- 14
- 15 46 Berlin JE, Storti KL, Brach JS. Using Activity Monitors to Measure Physical Activity in Free-Living
- 16 Conditions. *Physical Therapy* 2006;**86**:1137–45. doi:10.1093/ptj/86.8.1137
- 17
- 18 47 Strath SJ, Kaminsky LA, Ainsworth BE, *et al*. Guide to the assessment of physical activity: Clinical
- 19 and research applications: a scientific statement from the American Heart Association. *Circulation*
- 20 2013;**128**:2259–79. doi:10.1161/01.cir.0000435708.67487.da
- 21
- 22 48 Morris S, Earl K. Health Survey for England 2016 Well-being and mental health. *Health and Social*
- 23 *Care Information Centre* 2017.
- 24
- 25 49 Wade M, Mann S, Copeland RJ, *et al*. The effect of exercise referral schemes upon health and
- 26 wellbeing: Initial observational insights using individual patient data meta-analysis from The National

1
2
3 1 Referral database. doi:10.31236/osf.io/yebmr
4
5 2
6
7 3 50 Shah N, Cader M, Andrews WP, *et al.* Responsiveness of the Short Warwick Edinburgh Mental
8
9
10 4 Well-Being Scale (SWEMWBS): evaluation a clinical sample. *Health and Quality of Life Outcomes*
11
12 5 2018;**16**:239. doi:10.1186/s12955-018-1060-2
13
14 6
15
16 7 51 Cavill N, Roberts K, Rutter H. Standard evaluation framework for physical activity interventions.
17
18 8 Oxford: National Obesity Observatory 2012.
19
20 9
21
22
23 10 52 Craig P, Dieppe P, Macintyre S, *et al.* Developing and evaluating complex interventions: the new
24
25 11 Medical Research Council guidance. *BMJ* 2008;**337**:a1655. doi:10.1136/bmj.a1655
26
27 12
28
29
30 13 53 Rutter H, Savona N, Glonti K, *et al.* The need for a complex systems model of evidence for public
31
32 14 health. *Lancet* 2017;**17**:1267–9. doi:10.1016/S0140-6736
33
34 15
35
36 16 54 Beedie C, Mann S, Jimenez A, *et al.* Death by effectiveness: exercise as medicine caught in the
37
38 17 efficacy trap! *Brit J Sport Med* 2015;**0**:1–2. doi:10.1136/bjsports-2014-094389
39
40 18
41
42
43 19 55 Perera R, Heneghan C, Yudkin P. Graphical method for depicting randomised trials of complex
44
45 20 interventions. *BMJ* 2007;**334**:127. doi:10.1136/bmj.39045.396817.68
46
47
48
49 21
50
51
52
53
54
55
56
57
58
59
60







Figure 1. PaT Plot' describing intervention arm components.



TREND Statement Checklist

Paper Section/Topic	Item No.	Descriptor	Reported?	
			✓	Pg #
TITLE and ABSTRACT				
Title and Abstract	1	• Information on how units were allocated to interventions	✓	1,2
		• Structured abstract recommended	✓	2
		• Information on target population or study sample	✓	2
INTRODUCTION				
Background	2	• Scientific background and explanation of rationale	✓	4-5
		• Theories used in designing behavioral interventions	✓	6
METHODS				
Participants	3	• Eligibility criteria for participants, including criteria at different levels in recruitment/sampling plan (e.g., cities, clinics, subjects)	✓	5-6
		• Method of recruitment (e.g., referral, self-selection), including the sampling method if a systematic sampling plan was implemented	✓	5-6
		• Recruitment setting	✓	6,7
		• Settings and locations where the data were collected	✓	5-6
Interventions	4	• Details of the interventions intended for each study condition and how and when they were actually administered, specifically including:	✓	6-8
		◦ Content: what was given?	✓	6-8
		◦ Delivery method: how was the content given?	✓	6-8
		◦ Unit of delivery: how were subjects grouped during delivery?	✓	6-8
		◦ Deliverer: who delivered the intervention?	✓	6-8
		◦ Setting: where was the intervention delivered?	✓	6-8
		◦ Exposure quantity and duration: how many sessions or episodes or events were intended to be delivered? How long were they intended to last?	✓	6-8
		◦ Time span: how long was it intended to take to deliver the intervention to each unit?	✓	6-8
	◦ Activities to increase compliance or adherence (e.g., incentives)		N/A	
Objectives	5	• Specific objectives and hypotheses	✓	5
Outcomes	6	• Clearly defined primary and secondary outcome measures	✓	7-8
		• Methods used to collect data and any methods used to enhance the quality of measurements	✓	7-8
		• Information on validated instruments such as psychometric and biometric properties		N/A
Sample size	7	• How sample size was determined and, when applicable, explanation of any interim analyses and stopping rules	✓	8
Assignment method	8	• Unit of assignment (the unit being assigned to study condition, e.g., individual, group, community)		N/A
		• Method used to assign units to study conditions, including details of any restriction (e.g., blocking, stratification, minimization)		N/A
		• Inclusion of aspects employed to help minimize potential bias induced due to non-randomization (e.g., matching)		N/A
Blinding (masking)	9	• Whether or not participants, those administering the interventions, and those assessing the outcomes were blinded to study condition assignment; if so, statement regarding how the blinding was accomplished and how it was assessed		N/A
Unit of Analysis	10	• Description of the smallest unit that is being analysed to assess intervention effects (e.g., individual, group, or community)	✓	8-9
		• If the unit of analysis differs from the unit of assignment, the analytical method used to account for this (e.g., adjusting the standard error estimates by the design effect or using multilevel analysis)		N/A
Statistical methods	11	• Statistical methods used to compare study groups for primary methods outcome(s), including complex methods for correlated data	✓	8-9
		• Statistical methods used for additional analyses, such as subgroup analyses and adjusted analysis	✓	8-9
		• Methods for imputing missing data, if used		N/A

TREND Statement Checklist

		• Statistical software or programs used	✓	8-9
RESULTS				
Participant flow	12	<ul style="list-style-type: none"> Flow of participants through each stage of the study: enrollment, assignment, allocation and intervention exposure, follow-up, analysis (a diagram is strongly recommended) Enrollment: the numbers of participants screened for eligibility, found to be eligible or not eligible, declined to be enrolled, and enrolled in the study Assignment: the numbers of participants assigned to a study condition Allocation and intervention exposure: the number of participants assigned to each study condition and the number of participants who received each intervention Follow-up: the number of participants who completed the follow-up or did not complete the follow-up (i.e., lost to follow-up), by study condition Analysis: the number of participants included in or excluded from the main analysis, by study condition Description of protocol deviations from study as planned, along with reasons 	✓	9
Recruitment	13	• Dates defining the periods of recruitment and follow-up	✓	9
Baseline data	14	<ul style="list-style-type: none"> Baseline demographic and clinical characteristics of participants in each study condition Baseline characteristics for each study condition relevant to specific disease prevention research Baseline comparisons of those lost to follow-up and those retained, overall and by study condition Comparison between study population at baseline and target population of interest 	✓	10
Baseline equivalence	15	• Data on study group equivalence at baseline and statistical methods used to control for baseline differences		N/A
Numbers analyzed	16	<ul style="list-style-type: none"> Number of participants (denominator) included in each analysis for each study condition, particularly when the denominators change for different outcomes; statement of the results in absolute numbers when feasible Indication of whether the analysis strategy was "intention to treat" or, if not, description of how non-compliers were treated in the analyses 	✓	10-13
Outcomes and estimation	17	<ul style="list-style-type: none"> For each primary and secondary outcome, a summary of results for each estimation study condition, and the estimated effect size and a confidence interval to indicate the precision Inclusion of null and negative findings Inclusion of results from testing pre-specified causal pathways through which the intervention was intended to operate, if any 	✓	10
Ancillary analyses	18	• Summary of other analyses performed, including subgroup or restricted analyses, indicating which are pre-specified or exploratory		N/A
Adverse events	19	• Summary of all important adverse events or unintended effects in each study condition (including summary measures, effect size estimates, and confidence intervals)	✓	10
DISCUSSION				
Interpretation	20	<ul style="list-style-type: none"> Interpretation of the results, taking into account study hypotheses, sources of potential bias, imprecision of measures, multiplicative analyses, and other limitations or weaknesses of the study Discussion of results taking into account the mechanism by which the intervention was intended to work (causal pathways) or alternative mechanisms or explanations Discussion of the success of and barriers to implementing the intervention, fidelity of implementation Discussion of research, programmatic, or policy implications 	✓	14-17
Generalizability	21	• Generalizability (external validity) of the trial findings, taking into account the study population, the characteristics of the intervention, length of follow-up, incentives, compliance rates, specific sites/settings involved in the study, and other contextual issues	✓	14-17
Overall evidence	22	• General interpretation of the results in the context of current evidence and current theory	✓	14-17

From: Des Jarlais, D. C., Lyles, C., Crepaz, N., & the Trend Group (2004). Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: The TREND statement. *American Journal of Public Health*, 94, 361-366. For more information, visit: <http://www.cdc.gov/trendstatement/>